

**TABLE 3B-4
RATES^a OF REPORTED CASES OF GONORRHEA, CHLAMYDIA, AND EARLY SYPHILIS
BY AGE AND GENDER, ARIZONA, 2014**

Age group	GONORRHEA			CHLAMYDIA			EARLY SYPHILIS		
	Males	Females	Total	Males	Females	Total	Males	Females	Total
0-4	0.0	**	**	0.0	0.0	0.0	0.0	0.0	0.0
5-9	0.0	0.0	0.0	0.0	**	**	0.0	0.0	0.0
10-14	**	7.1	4.6	10.3	63.7	36.5	0.0	0.0	0.0
15-19	215.5	306.9	260.0	687.9	2656.2	1645.6	15.4	4.1	9.9
20-24	459.5	507.6	482.7	1364.8	3762.0	2521.5	63.6	8.5	37.0
25-29	381.1	319.5	351.4	872.6	1794.0	1315.9	55.8	11.5	34.5
30-34	240.6	198.3	219.9	459.3	842.7	646.3	49.5	7.4	29.0
35-39	159.7	105.8	133.0	253.0	417.0	334.4	35.4	4.9	20.2
40-44	106.3	60.5	83.5	159.7	191.6	175.6	35.4	4.3	19.9
45-49	96.2	31.5	63.9	94.7	91.1	92.9	43.2	**	21.9
50-54	52.6	16.4	34.2	51.2	47.8	49.5	28.2	**	14.1
55-59	25.7	7.9	16.4	33.8	28.2	30.9	14.6	**	7.2
60-64	20.7	**	10.3	16.8	8.0	12.1	7.9	0.0	3.7
65-over	5.9	**	3.0	5.3	2.8	3.9	1.4	0.0	0.7
Total	122.6	102.9	112.7	283.2	657.7	471.7	23.6	2.7	13.1

Notes: ** Cell suppressed due to rate/ratio/percent based on non-zero count less than 6;^a Number of cases per 100,000 population; table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2014; denominators for unknown or transgender category are not available; rates per 100,000 population.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.