

TABLE 3B-5
FREQUENCY OF REPORTED CASES, PERCENT DISTRIBUTION AND RATES OF EARLY AND LATE SYPHILIS,
GONORRHEA, AND CHLAMYDIA BY RACE/ETHNICITY, ARIZONA, 2014

Race/ethnicity	SYPHILIS						GONORRHEA						CHLAMYDIA		
	Early			Late			Resistant			Total					
	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate
White Non-Hispanic	350	39.8	9.1	125	22.6	3.2	0	0.0	0.0	1,564	20.6	40.6	5,203	16.4	135.2
Black or African American	90	10.2	29.3	62	11.2	20.2	0	0.0	0.0	1,034	13.6	336.2	2,357	7.4	766.3
Hispanic or Latino	332	37.7	16.1	232	41.9	11.3	0	0.0	0.0	2,135	28.1	103.8	9,282	29.2	451.4
Asian or Pacific Islander	13	1.5	5.6	8	1.4	3.4	0	0.0	0.0	58	0.8	24.8	268	0.8	114.7
American Indian or Alaska Native	48	5.5	16.9	24	4.3	8.4	0	0.0	0.0	587	7.7	206.5	2,804	8.8	986.6
Not Specified	47	5.3	N/A	103	18.6	N/A	*	**	N/A	2,207	29.1	N/A	11,836	37.3	N/A
Total	880	100.0	13.1	554	100.0	8.2	*	**	0.0	7,585	100.0	112.7	31,750	100.0	471.7

Notes: * Cell suppressed due to non-zero count less than 6; ** Cell suppressed due to rate/ratio/percent based on non-zero count less than 6; Table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2014; rates per 100,000 population.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.