

**TABLE 3B-5
 FREQUENCY OF REPORTED CASES, PERCENT DISTRIBUTION AND RATES OF EARLY AND LATE SYPHILIS,
 GONORRHEA, CHLAMYDIA AND HERPES BY ETHNICITY, ARIZONA, 2000**

Race	SYPHILIS						GONORRHEA						CHLAMYDIA			HERPES		
	Early			Late			Resistant			Total			Cases	%	Rate	Cases	%	Rate
Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases						
White	103	23.6	3.2	65	16.9	2.0	0	0	0	967	23.6	29.5	3,166	25.3	96.7	518	46.3	15.8
Black	96	22.0	60.4	69	17.9	43.4	0	0	0	1,003	24.4	630.8	1,091	8.7	686.2	76	6.8	47.8
Hispanic	200	45.8	15.4	204	53.0	15.8	0	0	0	1,449	35.3	111.8	4,540	36.3	350.3	221	19.8	17.1
Asian	3	0.7	3.0	1	0.3	1.1	0	0	0	22	0.5	22.2	89	0.7	89.9	7	0.7	7.1
American Indian	31	7.1	12.1	31	8.1	12.1	0	0	0	269	6.6	105.1	2,161	17.3	844.1	86	7.7	33.6
Not Specified	4	0.9	NA	15	3.9	NA	0	0	NA	395	9.6	NA	1,468	11.7	NA	211	18.9	NA
Total	437	100	8.5	385	100	7.5	0	100	0	4,105	100	80.0	12,515	100	243.9	1,119	100	21.8

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV/STD Services