

**TABLE 3B-5
FREQUENCY OF REPORTED CASES, PERCENT DISTRIBUTION AND RATES OF EARLY AND LATE SYPHILIS,
GONORRHEA, CHLAMYDIA AND HERPES BY ETHNICITY, ARIZONA, 2001**

Race	SYPHILIS						GONORRHEA						CHLAMYDIA			HERPES					
	Early			Late			Resistant			Total			Cases	%	Rate	Cases	%	Rate	Cases	%	Rate
	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate									
White	87	19.5	2.6	98	14.7	2.9	1	20.0	*	1,097	28.0	32.4	3,947	27.5	116.5	599	51.1	17.7			
Black	86	19.3	52.4	88	13.2	53.7	1	20.0	0.6	916	23.4	558.5	1,241	8.7	756.7	71	6.1	43.3			
Hispanic	223	50.0	16.6	386	57.9	28.8	2	40.0	0.2	1,336	34.1	99.7	6,250	43.6	466.4	319	27.2	23.8			
Asian	0	0	0	6	0.9	5.9	0	0	0	11	0.3	10.8	116	0.8	113.7	14	1.2	13.7			
American Indian	40	9.0	15.1	29	4.4	10.9	1	20.0	0	306	7.8	115.5	1,958	13.6	738.9	74	6.3	27.9			
Not Specified	10	2.2	NA	60	0.9	NA	0	0	NA	256	6.5	NA	840	5.9	NA	96	8.2	NA			
Total	446	100%	8.4	667	100%	12.6	5	100%	0.1	3,922	100%	73.9	14,352	100%	270.4	1,173	100%	22.1			

* Quantity equal or less than 0.05

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV/STD Services