

**TABLE 4
MONITORING PROGRESS TOWARD ARIZONA AND SELECTED NATIONAL YEAR 2010 OBJECTIVES**

Focus areas and selected objectives: (in parentheses are <i>Healthy People 2010</i> objective numbers)	Baseline for the U.S.	ARIZONA RATES, RATIOS OR CASES BY YEAR:										2010 TARGET		
		2000	2001	2002	2003	2004	2005	2006	2007	2008	2009		2010	
4A. MATERNAL, INFANT, AND CHILD HEALTH														
Reduce fetal deaths at 20 or more weeks of gestation (HP16-1a)	6.8	7.6	8.2	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	4.1
Reduce fetal and infant deaths during perinatal period (HP16-1b)	7.5	5.3	6.6	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	4.5
♦ Reduce infant deaths (HP16-1c)	7.2	6.7	6.9	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	♦6.0
Reduce neonatal deaths (HP16-1d)	4.8	4.3	4.5	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	2.9
Reduce postneonatal deaths (HP16-1e)	2.4	2.4	2.4	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	1.2
Reduce infant deaths due to birth defects (HP16-1f)	1.6	1.6	1.7	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	1.1
Reduce deaths from sudden infant death syndrome (SIDS) (HP16-1h)	0.7	0.3	0.4	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	0.25
Reduce the rate of death among children aged 1 to 4 years (HP16-2a)	34.6	36.7	46.0	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	18.6
Reduce the rate of death among children aged 5 to 9 years (HP16-2b)	17.7	19.2	20.1	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	12.3
Reduce the rate of death among children aged 10 to 14 years (HP16-3a)	22.1	22.2	24.0	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	16.8
Reduce the rate of death among adolescents aged 15 to 19 years (HP16-3b)	70.6	78.9	86.6	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	39.8
Reduce the rate of death among young adults aged 20 to 24 years (HP16-3c)	95.3	104.7	106.3	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	49.0
Reduce maternal deaths (HP16-4)	7.1	*8.2	*9.4	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	3.3
Increase the proportion of pregnant women who receive prenatal care in the first trimester (HP16-6a)	85%	75%	75.5%	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	90%
Reduce low birth weight (LBW) (HP16-10a)	7.6%	7.2%	7.0%	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	5.0%
Reduce very low birth weight (VLBW) (HP16-10b)	1.4%	1.3%	1.1%	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	0.9%
Reduce preterm births (HP16-11a)	11.6%	10.3%	9.9%	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	7.6%
Increase abstinence from cigarette smoking among pregnant women (HP16-17c)	87.0%	92.6%	93.2%	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	99.0%

♦ Indicates objectives and targets identified in **Healthy Arizona 2010**.

Note: fetal and perinatal deaths are per 1,000 live births+fetal deaths. Perinatal period= 28 weeks of gestation to 7 days after birth. Infant, neonatal, and postneonatal deaths are per 1,000 live births. Infant deaths due to birth defects, congenital birth defects, and SIDS are per 1,000 live births. All age-specific mortality rates are per 100,000 persons. The maternal mortality ratio is per 100,000 live births. All other proportions and ratios are per 100 live births. Preterm births= births prior to 37 completed weeks of gestation. The baseline year for the U.S. is 1997.

* These rates are based on fewer than 10 cases and are not statistically reliable.