

**TABLE 5E-28
RATES* FOR THE LEADING CAUSES OF DEATH AMONG ADOLESCENTS 15-19 YEARS
BY COUNTY OF RESIDENCE, ARIZONA, 2001**

| | Arizona | Apache | Cochise | Coconino | Gila | Graham | Greenlee | Maricopa | Mohave | Navajo | Pima | Pinal | Santa Cruz | Yavapai | Yuma | La Paz |
|---|---------|--------|---------|----------|-------|--------|----------|----------|--------|--------|------|-------|------------|---------|------|--------|
| Total, all causes | 86.6 | 218.7 | 64.9 | 77.5 | 204.7 | 62.1 | .0 | 77.5 | 135.8 | 193.5 | 80.2 | 94.6 | 30.5 | 117.5 | 48.6 | 83.5 |
| Accident (unintentional injury) | 37.6 | 102.0 | .0 | 38.7 | 117.0 | 31.0 | .0 | 33.1 | 41.8 | 86.0 | 34.6 | 39.4 | 30.5 | 81.4 | 24.3 | 83.5 |
| Transport accidents | 28.4 | 72.9 | .0 | 29.0 | 87.7 | 31.0 | .0 | 25.1 | 31.3 | 53.8 | 22.0 | 39.4 | 30.5 | 81.4 | 16.2 | 83.5 |
| Motor vehicle accidents | 26.3 | 58.3 | .0 | 29.0 | 58.5 | 31.0 | .0 | 23.7 | 20.9 | 53.8 | 20.4 | 39.4 | 30.5 | 72.3 | 16.2 | 83.5 |
| Nontransport accidents | 9.2 | 29.2 | .0 | 9.7 | 29.2 | .0 | .0 | 8.1 | 10.4 | 32.3 | 12.6 | .0 | .0 | .0 | 8.1 | .0 |
| Drowning and submersion | 1.3 | .0 | .0 | .0 | .0 | .0 | .0 | 2.2 | .0 | .0 | .0 | .0 | .0 | .0 | .0 | .0 |
| Exposure to smoke, fire and flames | 1.1 | .0 | .0 | .0 | 29.2 | .0 | .0 | .0 | .0 | 32.3 | .0 | .0 | .0 | .0 | .0 | .0 |
| Accidental discharge of firearms | 1.6 | .0 | .0 | .0 | .0 | .0 | .0 | 1.8 | .0 | .0 | 3.1 | .0 | .0 | .0 | .0 | .0 |
| Accidental poisoning | 2.6 | .0 | .0 | 9.7 | .0 | .0 | .0 | 1.8 | .0 | .0 | 7.9 | .0 | .0 | .0 | .0 | .0 |
| Assault (homicide) | 12.9 | 14.6 | .0 | .0 | .0 | 31.0 | .0 | 15.7 | 10.4 | .0 | 14.1 | 7.9 | .0 | 9.0 | .0 | .0 |
| Intentional self-harm (suicide) | 10.5 | 29.2 | 32.4 | 29.0 | 29.2 | .0 | .0 | 6.7 | 10.4 | 75.3 | 7.9 | 23.6 | .0 | .0 | .0 | .0 |
| Malignant neoplasms | 5.3 | .0 | 10.8 | .0 | .0 | .0 | .0 | 4.9 | 10.4 | 10.8 | 4.7 | .0 | .0 | 9.0 | 16.2 | .0 |
| Diseases of heart | 2.6 | .0 | .0 | .0 | .0 | .0 | .0 | 3.1 | 20.9 | .0 | .0 | 7.9 | .0 | .0 | .0 | .0 |
| Congenital malformations | 1.3 | .0 | .0 | .0 | .0 | .0 | .0 | 1.8 | .0 | .0 | 1.6 | .0 | .0 | .0 | .0 | .0 |

* Number of deaths per 100,000 adolescents in specified county.
Note: rates based on fewer than 10 deaths are not statistically reliable (see Table 5E-27).