

**TABLE 5E-30
RATES* FOR THE LEADING CAUSES OF DEATH AMONG YOUNG ADULTS 20-44 YEARS
BY COUNTY OF RESIDENCE, ARIZONA, 2001**

	Arizona	Apache	Cochise	Coconino	Gila	Graham	Greenlee	Maricopa	Mohave	Navajo	Pima	Pinal	Santa Cruz	Yavapai	Yuma	La Paz
Total, all causes	158.2	401.9	156.4	138.1	243.2	145.9	109.3	145.2	217.4	321.7	156.4	188.6	134.2	195.3	118.7	308.6
Accident (unintentional injury)	48.1	116.8	65.2	48.9	66.3	42.9	.0	43.0	49.3	160.9	45.7	50.7	47.4	60.9	40.8	205.7
Transport accidents	25.4	107.5	41.7	36.1	36.8	34.3	.0	22.2	22.4	99.7	17.7	33.3	23.7	35.7	16.7	144.0
Motor vehicle accidents	22.9	107.5	36.5	29.7	29.5	25.7	.0	19.9	22.4	90.1	16.1	31.7	23.7	27.3	16.7	123.4
Nontransport accidents	22.6	9.3	23.5	12.7	29.5	8.6	.0	20.8	26.9	61.1	28.0	17.4	23.7	25.2	24.1	61.7
Accidental poisoning	14.1	.0	15.6	6.4	7.4	8.6	.0	13.2	11.2	12.9	21.6	12.7	7.9	16.8	13.0	.0
Malignant neoplasms	18.5	23.4	7.8	12.7	14.7	.0	.0	18.0	20.2	22.5	19.0	22.2	31.6	23.1	26.0	41.1
Breast	5.2	18.5	.0	4.2	.0	.0	.0	5.9	8.9	.0	2.6	3.8	.0	4.2	7.8	42.8
Trachea, bronchus and lung	1.8	.0	.0	.0	.0	.0	.0	1.9	4.5	.0	1.3	.0	7.9	6.3	.0	.0
Colon, rectum and anus	1.4	.0	.0	2.1	7.4	.0	.0	.8	6.7	3.2	2.6	1.6	7.9	2.1	1.9	.0
Assault (homicide)	16.3	9.3	13.0	12.7	36.8	.0	.0	19.3	13.4	19.3	9.3	17.4	15.8	6.3	5.6	.0
Intentional self-harm (suicide)	13.3	18.7	15.6	17.0	22.1	17.2	109.3	11.8	17.9	22.5	15.5	14.3	15.8	21.0	1.9	.0
Diseases of heart	12.2	9.3	10.4	8.5	.0	17.2	.0	12.1	22.4	6.4	12.6	14.3	7.9	10.5	13.0	41.1
Chronic liver disease and cirrhosis	5.4	18.7	5.2	6.4	14.7	.0	.0	4.0	9.0	35.4	5.2	9.5	.0	12.6	.0	.0
Human immunodeficiency virus (HIV) disease	4.6	4.7	2.6	2.1	.0	.0	.0	5.0	2.2	.0	6.4	3.2	.0	4.2	.0	.0
Cerebrovascular diseases	2.4	4.7	.0	2.1	.0	.0	.0	2.2	2.2	3.2	2.3	9.5	.0	2.1	1.9	.0
Diabetes	2.2	4.7	5.2	.0	7.4	8.6	.0	1.8	4.5	.0	2.6	3.2	.0	2.1	3.7	.0
Influenza and pneumonia	1.8	4.7	2.6	4.2	14.7	.0	.0	1.8	4.5	.0	1.3	1.6	.0	.0	.0	.0

* Number of deaths per 100,000 persons 20-44 in specified county. The rate from breast cancer is per 100,000 females.
Note: rates based on fewer than 10 deaths are not statistically reliable (see Table 5E-29).