

**TABLE 2D-4**  
**PERCENT DEATHS BEFORE EXPECTED\* YEARS OF LIFE REACHED FOR THE LEADING CAUSES OF MORTALITY BY GENDER, RACE/ETHNICITY AND URBAN/RURAL AREAS\*, ARIZONA, 2002**

	TOTAL STATE			AREA/GENDER:						RACE/ETHNICITY:				
	TOTAL	MALE	FEMALE	TOTAL URBAN	URBAN MALE	URBAN FEMALE	TOTAL RURAL	RURAL MALE	RURAL FEMALE	WHITE NON-HISPANIC	HISPANIC	BLACK	AMERICAN INDIAN	ASIAN
<b>Total, all causes</b>	50.2%	58.4%	41.1%	49.2%	57.7%	40.0%	54.1%	60.9%	45.5%	45.6%	70.3%	71.7%	78.3%	59.8%
<b>Major cardiovascular diseases</b>	37.7%	47.8%	27.1%	36.3%	46.6%	26.0%	43.5%	52.4%	32.6%	34.8%	54.5%	58.0%	64.7%	42.9%
<b>Diseases of heart</b>	40.1%	50.3%	28.4%	38.5%	48.9%	27.1%	47.3%	56.1%	34.7%	37.5%	55.7%	61.8%	64.9%	43.3%
<b>Malignant neoplasms</b>	62.0%	63.5%	60.2%	61.3%	62.8%	59.5%	64.8%	66.0%	63.4%	60.1%	71.7%	78.6%	74.1%	76.1%
<b>Lung cancer</b>	66.7%	69.3%	63.4%	65.7%	68.0%	62.9%	70.5%	74.4%	65.5%	65.7%	73.6%	88.3%	83.3%	40.0%
<b>Prostate cancer</b>	32.6%	32.6%	.0%	32.7%	32.7%	.0%	31.8%	31.8%	.0%	30.2%	44.6%	58.8%	40.0%	.0%
<b>Female breast cancer</b>	70.1%	60.0%	70.3%	70.9%	57.1%	71.1%	66.9%	66.7%	66.9%	66.5%	84.6%	91.7%	100.0%	100.0%
<b>Colorectal cancer</b>	56.6%	61.0%	51.4%	55.9%	59.9%	50.9%	59.9%	65.9%	53.2%	54.5%	71.1%	64.0%	76.9%	60.0%
<b>Cerebrovascular diseases</b>	28.6%	36.2%	23.3%	28.6%	36.6%	23.2%	28.3%	34.2%	23.9%	24.7%	51.1%	50.0%	57.1%	37.5%
<b>Chronic lower respiratory diseases</b>	42.1%	44.3%	39.8%	41.8%	44.4%	39.2%	43.3%	43.7%	42.9%	41.3%	52.3%	60.6%	46.2%	50.0%
<b>Accidents (unintentional injuries)</b>	78.7%	84.4%	67.8%	78.3%	84.6%	66.4%	80.1%	83.6%	72.7%	70.9%	93.9%	93.8%	95.3%	83.3%
<b>Motor vehicle accidents</b>	92.4%	94.0%	89.0%	92.2%	94.6%	87.2%	92.9%	92.4%	94.1%	87.2%	98.8%	100.0%	98.3%	87.5%
<b>Falls</b>	33.5%	41.0%	25.2%	33.2%	41.6%	24.3%	35.1%	38.6%	30.3%	28.4%	64.9%	60.0%	72.2%	.0%
<b>Accidental poisoning</b>	99.2%	99.4%	98.5%	99.5%	99.7%	99.1%	97.5%	98.2%	95.8%	98.8%	100.0%	100.0%	100.0%	.0%
<b>Accidental drowning</b>	89.9%	95.7%	77.3%	87.9%	94.4%	77.3%	100.0%	100.0%	.0%	86.4%	94.7%	100.0%	100.0%	100.0%
<b>Influenza and pneumonia</b>	29.8%	36.2%	24.5%	29.6%	35.4%	24.7%	31.0%	39.6%	23.9%	25.9%	51.6%	51.9%	44.3%	40.0%
<b>Alzheimer's disease</b>	9.4%	13.7%	7.4%	9.7%	14.8%	7.3%	7.8%	7.5%	7.9%	8.7%	14.9%	27.8%	22.2%	.0%
<b>Diabetes</b>	58.5%	62.8%	54.2%	57.9%	62.0%	53.8%	61.4%	66.1%	56.2%	53.5%	64.0%	73.7%	79.3%	60.0%
<b>Intentional self-harm (suicide)</b>	91.0%	90.8%	92.0%	91.4%	91.4%	91.6%	89.7%	88.9%	93.8%	89.3%	96.1%	100.0%	100.0%	100.0%
<b>Chronic liver disease and cirrhosis</b>	90.3%	92.9%	86.1%	89.0%	91.8%	84.7%	95.3%	97.5%	91.8%	86.8%	93.6%	100.0%	100.0%	100.0%
<b>Nephritis, nephrotic syndrome and nephrosis</b>	42.5%	45.6%	39.8%	40.6%	44.3%	37.5%	49.3%	49.3%	49.2%	34.5%	60.2%	66.7%	77.8%	50.0%
<b>Septicemia</b>	56.4%	61.1%	51.6%	55.9%	61.8%	49.7%	58.2%	58.3%	58.0%	49.7%	69.8%	57.1%	89.5%	.0%
<b>Assault (homicide)</b>	98.9%	99.7%	95.1%	99.0%	100.0%	93.9%	98.4%	98.0%	100.0%	98.0%	100.0%	97.4%	100.0%	50.0%
<b>Parkinson's disease</b>	21.7%	22.6%	20.0%	22.4%	23.2%	21.0%	17.7%	19.6%	12.5%	21.4%	25.0%	100.0%	33.3%	.0%
<b>Essential (primary) hypertension and hypertensive renal disease</b>	27.6%	33.8%	22.7%	26.8%	32.5%	22.2%	34.4%	46.2%	26.3%	21.8%	56.3%	33.3%	85.7%	50.0%
<b>Human immunodeficiency virus (HIV) disease</b>	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	.0%	100.0%	100.0%	100.0%	100.0%	.0%
<b>INJURY BY FIREARMS</b>	92.9%	92.7%	93.8%	93.8%	93.8%	94.2%	88.6%	88.1%	92.0%	89.5%	98.8%	100.0%	100.0%	80.0%
<b>DRUG-INDUCED DEATHS</b>	97.8%	99.5%	94.4%	98.0%	99.7%	94.5%	96.9%	98.4%	93.8%	97.2%	99.1%	100.0%	100.0%	.0%
<b>ALCOHOL-INDUCED DEATHS</b>	94.9%	95.1%	94.2%	94.3%	93.9%	95.3%	96.7%	98.9%	91.4%	94.4%	91.4%	91.7%	100.0%	100.0%

\* Expected years of life at birth for all U.S. residents.

\*\* Urban = Maricopa, Pima, Pinal and Yuma counties.

Note: The causes of death for 2002 are classified by the Tenth Revision of the International Classification of Diseases (ICD-10).