

**TABLE 3B-5
 FREQUENCY OF REPORTED CASES, PERCENT DISTRIBUTION AND RATES OF EARLY AND LATE SYPHILIS, GONORRHEA,
 CHLAMYDIA AND HERPES BY ETHNICITY, ARIZONA, 2002**

Race/Ethnicity	SYPHILIS						GONORRHEA						CHLAMYDIA			HERPES		
	Early			Late			Resistant			Total			Cases	%	Rate	Cases	%	Rate
	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate						
White	88	22	2.5	105	15.5	3.0	0	0	0.0	1,064	28.2	30.5	3,955	26.6	113.2	566	49.3	16.2
Black	68	17	40.2	88	13.0	52.1	1	100	0.6	730	19.4	432.0	1,183	7.9	700.0	76	6.6	45.0
Hispanic	174	43.5	12.6	396	58.6	28.7	0	0	0.0	1,363	36.1	98.6	6,265	42.1	453.3	272	23.7	19.7
Asian	1	0.3	0.9	13	1.9	12.3	0	0	0.0	12	0.3	11.3	111	0.8	104.7	8	0.7	7.6
American Indian	44	11	16.1	38	5.6	13.9	0	0	0.0	294	7.8	107.7	1,957	13.1	716.9	98	8.5	35.9
Not Specified	25	6.2	NA	37	5.5	NA	0	0	0.0	309	8.2	NA	1,428	9.6	NA	128	11.2	NA
Total	400	100	7.3	677	100	12.4	1	100	0.0	3,772	100	68.9	14,899	100	272.2	1,148	100	21.0

* Quantity equal or less than 0.05

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV/STD Services