

**TABLE 5E-22
POSTNEONATAL MORTALITY RATES¹ BY COUNTY OF RESIDENCE,
ARIZONA AND UNITED STATES, 1992-2002**

| County | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 |
|-------------------|------|------|------|------|------|------|------|------|------|------|------|
| U.S. | 3.1 | 3.0 | 3.0 | 2.7 | 2.5 | 2.5 | 2.4 | 2.4 | 2.3 | 2.3 | NA |
| ARIZONA | 3.6 | 2.8 | 3.2 | 2.8 | 2.7 | 2.7 | 2.8 | 2.3 | 2.4 | 2.4 | 2.2 |
| Apache | 4.9 | 2.5 | 4.8 | 6.0 | 2.3 | 3.8 | 1.6 | 4.5 | 4.7 | 4.7 | 1.8 |
| Cochise | 3.4 | 3.6 | 5.9 | 6.3 | 1.7 | 1.8 | 3.1 | 1.2 | 4.0 | 3.1 | 4.7 |
| Coconino | 2.7 | 3.8 | 4.0 | 3.4 | 4.9 | 3.9 | 6.3 | 0.0 | 2.7 | 2.7 | 1.6 |
| Gila | 5.7 | 1.5 | 5.9 | 1.5 | 3.1 | 1.5 | 1.4 | 6.0 | 5.9 | 4.6 | 2.9 |
| Graham | 0.0 | 2.2 | 4.8 | 2.5 | 4.3 | 2.0 | 2.0 | 0.0 | 2.2 | 4.5 | 2.1 |
| Greenlee | 0.0 | 12.5 | 0.0 | 0.0 | 0.0 | 0.0 | 7.1 | 0.0 | 0.0 | 0.0 | 0.0 |
| Maricopa | 3.6 | 2.6 | 2.9 | 2.6 | 2.5 | 2.5 | 2.4 | 2.2 | 2.2 | 2.1 | 2.1 |
| Mohave | 5.3 | 1.1 | 1.1 | 3.3 | 5.0 | 3.4 | 6.0 | 2.9 | 2.8 | 1.6 | 1.0 |
| Navajo | 5.4 | 1.1 | 3.9 | 4.2 | 4.1 | 3.0 | 6.2 | 2.3 | 1.2 | 3.8 | 3.7 |
| Pima | 3.0 | 3.4 | 3.0 | 2.7 | 2.0 | 2.2 | 2.6 | 2.9 | 2.6 | 2.4 | 3.0 |
| Pinal | 5.9 | 1.9 | 4.9 | 2.0 | 2.4 | 6.1 | 3.6 | 2.5 | 3.1 | 4.3 | 1.5 |
| Santa Cruz | 1.1 | 2.5 | 0.0 | 0.0 | 2.5 | 0.0 | 1.3 | 0.0 | 0.0 | 0.0 | 2.6 |
| Yavapai | 4.6 | 3.0 | 2.2 | 3.3 | 3.2 | 1.3 | 4.1 | 2.5 | 4.0 | 2.9 | 2.1 |
| Yuma | 1.4 | 2.4 | 3.3 | 0.7 | 1.4 | 2.1 | 2.8 | 1.4 | 2.0 | 3.3 | .7 |
| La Paz | 0.0 | 4.5 | 4.5 | 0.0 | 0.0 | 5.0 | 5.9 | 7.4 | 5.4 | 5.4 | 0.0 |

¹ Post-neonatal deaths per 1,000 live births.