

**TABLE 2D-4
PERCENT DEATHS BEFORE EXPECTED* YEARS OF LIFE REACHED FOR THE LEADING CAUSES OF MORTALITY BY GENDER,
RACE/ETHNICITY AND URBAN/RURAL AREAS*, ARIZONA, 2003**

	TOTAL STATE			AREA/GENDER:						RACE/ETHNICITY:				
	TOTAL	MALE	FEMALE	TOTAL URBAN	URBAN MALE	URBAN FEMALE	TOTAL RURAL	RURAL MALE	RURAL FEMALE	WHITE NON-HISPANIC	HISPANIC or LATINO	BLACK or AFRICAN-AMERICAN	AMERICAN INDIAN or ALASKA NATIVE	ASIAN or PACIFIC ISLANDER
Total, all causes	50.8%	59.0%	41.7%	49.8%	58.0%	40.9%	54.5%	62.6%	44.9%	46.1%	70.0%	70.6%	78.4%	65.7%
Major cardiovascular diseases	38.2%	48.2%	27.8%	36.7%	46.3%	27.0%	44.4%	55.5%	31.4%	35.3%	51.9%	58.3%	65.6%	61.0%
Diseases of heart	40.0%	50.1%	28.6%	38.1%	47.7%	27.4%	47.8%	59.1%	33.6%	37.3%	52.9%	61.2%	67.0%	57.1%
Coronary heart disease	40.4%	50.5%	28.3%	38.1%	47.9%	26.8%	50.9%	61.3%	35.7%	38.0%	52.8%	58.8%	66.2%	59.0%
Malignant neoplasms	62.3%	63.2%	61.2%	61.6%	62.5%	60.7%	64.7%	66.0%	63.1%	60.6%	71.8%	73.3%	69.2%	72.6%
Lung cancer	67.5%	69.0%	65.5%	66.3%	68.2%	63.9%	72.0%	72.3%	71.6%	67.3%	67.1%	71.7%	72.2%	66.7%
Prostate cancer	32.2%	32.2%	.0%	29.3%	29.3%	.0%	41.7%	41.7%	.0%	30.6%	38.5%	57.1%	50.0%	.0%
Female breast cancer	69.2%	60.0%	69.3%	70.3%	66.7%	70.3%	64.7%	50.0%	64.9%	66.9%	86.8%	70.8%	61.5%	87.5%
Colorectal cancer	56.9%	63.7%	48.9%	55.4%	62.5%	47.4%	63.5%	69.4%	56.3%	54.3%	69.4%	71.4%	80.0%	100.0%
Cerebrovascular diseases	31.0%	39.9%	24.9%	30.8%	39.8%	24.8%	32.0%	40.4%	25.3%	27.0%	46.3%	43.3%	68.0%	78.3%
Chronic lower respiratory diseases	43.8%	45.8%	41.8%	43.0%	45.2%	40.9%	47.1%	48.3%	45.6%	42.5%	57.3%	71.0%	64.0%	42.9%
Accidents (unintentional injuries)	78.7%	85.5%	65.9%	76.6%	83.5%	63.2%	85.2%	91.7%	73.5%	70.5%	93.5%	92.6%	95.9%	90.5%
Motor vehicle accidents	93.1%	95.2%	88.6%	91.8%	94.2%	86.4%	96.1%	97.6%	93.1%	89.3%	96.2%	96.2%	99.4%	100.0%
Falls	29.4%	37.9%	20.9%	26.8%	33.8%	19.8%	42.9%	57.5%	27.0%	25.5%	50.0%	50.0%	73.7%	75.0%
Accidental poisoning	98.9%	99.1%	98.6%	99.0%	99.0%	99.1%	98.5%	100.0%	96.9%	98.7%	100.0%	96.3%	100.0%	.0%
Accidental drowning	95.4%	94.4%	100.0%	94.6%	93.2%	100.0%	100.0%	100.0%	100.0%	91.5%	100.0%	100.0%	100.0%	100.0%
Influenza and pneumonia	33.5%	39.5%	26.9%	34.1%	40.1%	27.5%	31.6%	37.6%	25.5%	29.7%	48.5%	54.8%	47.4%	42.9%
Alzheimer's disease	10.8%	16.4%	8.3%	10.8%	16.0%	8.3%	11.1%	18.6%	8.1%	9.8%	22.8%	21.2%	16.7%	.0%
Diabetes	58.7%	64.8%	52.2%	58.9%	64.7%	52.5%	58.0%	65.3%	51.2%	52.4%	70.9%	75.4%	76.8%	42.9%
Intentional self-harm (suicide)	90.9%	89.8%	95.6%	90.3%	88.9%	95.8%	93.3%	92.9%	94.9%	88.8%	98.1%	100.0%	100.0%	83.3%
Chronic liver disease and cirrhosis	89.4%	92.9%	82.4%	89.3%	93.2%	80.8%	90.0%	91.8%	86.8%	86.7%	92.9%	100.0%	98.7%	.0%
Nephritis, nephrotic syndrome and nephrosis	45.8%	48.4%	43.3%	48.3%	51.0%	45.8%	37.9%	40.8%	34.4%	36.7%	62.2%	87.5%	78.4%	16.7%
Septicemia	57.1%	58.6%	55.7%	55.9%	58.1%	53.9%	60.9%	60.0%	61.8%	50.5%	69.6%	66.7%	81.0%	100.0%
Assault (homicide)	98.9%	99.5%	97.0%	98.8%	99.4%	96.4%	100.0%	100.0%	100.0%	97.0%	99.6%	100.0%	100.0%	100.0%
Parkinson's disease	22.4%	23.8%	20.6%	21.6%	24.4%	18.1%	26.7%	20.6%	34.6%	20.6%	33.3%	50.0%	62.5%	.0%
Essential (primary) hypertension and hypertensive renal disease	35.1%	38.3%	33.0%	37.2%	40.7%	35.0%	20.0%	20.0%	20.0%	28.2%	69.4%	73.3%	57.1%	25.0%
Human immunodeficiency virus (HIV) disease	99.4%	99.3%	100.0%	99.3%	99.3%	100.0%	100.0%	100.0%	100.0%	99.0%	100.0%	100.0%	100.0%	100.0%
INJURY BY FIREARMS	92.1%	91.6%	95.0%	92.1%	91.6%	94.9%	92.2%	91.7%	95.0%	86.9%	100.0%	100.0%	97.7%	100.0%
DRUG-INDUCED DEATHS	97.4%	98.4%	95.4%	97.3%	98.2%	95.3%	97.9%	100.0%	95.7%	97.0%	98.3%	97.3%	100.0%	.0%
ALCOHOL-INDUCED DEATHS	96.1%	96.4%	95.2%	96.2%	97.0%	93.3%	96.0%	94.8%	100.0%	95.0%	97.6%	100.0%	98.1%	.0%

* Expected years of life at birth for all U.S. residents.

** Urban = Maricopa, Pima, Pinal and Yuma counties.

Note: The causes of death for 2003 are classified by the Tenth Revision of the International Classification of Diseases (ICD-10).