

**TABLE 3B-5
 FREQUENCY OF REPORTED CASES, PERCENT DISTRIBUTION AND RATES OF EARLY AND LATE SYPHILIS, GONORRHEA,
 CHLAMYDIA AND HERPES BY ETHNICITY, ARIZONA, 2004**

	SYPHILIS						GONORRHEA						CHLAMYDIA			HERPES		
	Early			Late			Resistant			Total			Cases	%	Rate	Cases	%	Rate
Race/Ethnicity	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate						
White non-Hispanic	118	31	3.1	96	17	2.5	0	0	0.0	1,230	30	32.5	4,381	26	115.7	504	42	13.3
Black or African American	30	8	16.6	64	11	35.4	1	100	0.6	682	17	376.8	1,225	7	676.8	54	5	29.8
Hispanic or Latino	141	37	9.6	326	57	22.1	0	0	0.0	1,246	31	84.6	6,088	36	413.6	271	23	18.4
Asian or Pacific Islander	1	0	0.9	3	1	2.7	0	0	0.0	14	0	12.3	125	1	110.6	10	1	8.9
American Indian or Alaska Native	42	11	14.9	23	4	8.1	0	0	0.0	403	10	142.7	2,461	15	872.7	102	9	36.2
Not Specified	48	13	NA	65	10	NA	0	0	0.0	513	12	NA	2,589	15	NA	255	20	NA
Total	380	100	6.5	577	100	9.9	1	100	0.0	4,088	100	70.1	16,869	100	289.2	1,196	100	20.5

* Quantity equal or less than 0.05

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, OIDS, STD Control Program