

**TABLE 1C-4
SELECTED CHARACTERISTICS OF REPORTABLE SPONTANEOUS¹ FETAL LOSSES BY YEAR, ARIZONA, 2000-2005**

Characteristic:		YEAR					
		2000	2001	2002	2003	2004	2005
Total reportable spontaneous fetal losses		532	566	556	539	505	532
Gender	Male	280	275	299	296	231	262
	Female	244	267	253	240	270	268
	Unknown	8	24	4	2	4	1
Mother's race/ethnicity	White non-Hispanic	223	234	212	199	210	208
	Hispanic or Latino	231	228	256	246	214	232
	Black or African American	28	33	35	41	30	33
	American Indian or Alaska Native	29	53	37	41	31	45
	Asian or Pacific Islander	9	14	12	8	17	11
	Unknown	12	4	4	4	3	3
Mother's age group	<15	2	6	2	1	1	0
	15-17	42	28	24	31	25	29
	18-19	47	51	40	55	50	44
	20-24	125	125	146	134	130	141
	25-29	126	137	129	140	106	125
	30-34	92	115	111	93	105	112
	35-39	64	71	69	62	67	58
	40-44	25	20	27	17	17	17
	45+	2	1	4	1	1	2
	Unknown	7	12	4	5	3	4
Gestational age²	20-27 weeks	271	290	277	285	254	263
	28+ weeks	260	274	278	252	249	266
	Unknown	1	2	1	2	2	3
Weight in grams	<350 grams	82	74	98	90	83	95
	350-499	91	94	78	95	83	79
	500-2,499	236	214	221	243	219	225
	2,501-3,999	91	101	115	86	84	102
	4,000+	7	16	15	12	21	13
	Unknown	25	67	29	13	15	18

¹ Include spontaneous terminations of pregnancy at 20 or more weeks of gestation (or if gestational age is unknown, the deaths of fetuses of at least 350 grams in weight), exclude induced terminations of pregnancy.

²For statistical purposes, fetal deaths are classified according to gestational age. A death that occurs at 20 or more weeks of gestation constitutes a fetal death, and after 28 weeks it is considered a late fetal death.

Note: The number of reportable spontaneous fetal deaths for 2004 was revised. The values in the 2004 column above may differ from those previously published.