

TABLE 2D-4
PERCENT DEATHS BEFORE EXPECTED* YEARS OF LIFE REACHED FOR THE LEADING CAUSES OF MORTALITY BY
GENDER, RACE/ETHNICITY AND URBAN/RURAL AREAS, ARIZONA, 2005**

	TOTAL STATE			AREA/GENDER:						RACE/ETHNICITY:					
	Total	Male	Female	Total Urban	Urban Male	Urban Female	Total Rural	Rural Male	Rural Female	White Non-Hispanic	Hispanic or Latino	Black or African American	American Indian or Alaska Native	Asian or Pacific Islander	
Total, all causes	52.70%	60.80%	43.70%	51.80%	60.20%	42.70%	56.20%	63.20%	47.60%	47.90%	71.30%	75.10%	78.40%	65.10%	
Major cardiovascular diseases	40.20%	50.30%	29.70%	39.10%	49.10%	29.00%	44.80%	54.40%	32.70%	37.00%	55.80%	63.90%	68.10%	45.50%	
Diseases of heart	42.20%	52.10%	30.60%	40.80%	50.80%	29.80%	47.50%	56.90%	34.20%	39.00%	58.20%	64.80%	71.40%	49.30%	
Coronary heart disease	42.80%	53.10%	30.10%	40.90%	51.40%	28.80%	51.50%	59.90%	37.30%	40.10%	57.20%	63.90%	69.10%	49.10%	
Malignant neoplasms	64.00%	65.00%	62.80%	63.20%	64.70%	61.50%	66.80%	66.10%	67.70%	62.10%	72.60%	79.10%	76.30%	80.80%	
Lung cancer	68.20%	70.00%	66.00%	67.00%	69.70%	63.70%	72.80%	71.40%	74.60%	67.70%	68.80%	86.00%	66.70%	75.00%	
Prostate cancer	38.70%	38.70%	0.00%	36.80%	36.80%	0.00%	45.00%	45.00%	0.00%	37.00%	51.70%	44.40%	22.20%	100.00%	
Female breast cancer	69.90%	57.10%	70.00%	68.40%	50.00%	68.60%	75.20%	100.00%	75.00%	66.50%	86.70%	90.00%	84.60%	85.70%	
Colorectal cancer	58.00%	62.70%	52.70%	56.80%	61.80%	51.30%	63.10%	66.00%	59.20%	55.30%	72.70%	71.40%	81.80%	66.70%	
Malignant melanoma of skin	70.40%	71.40%	66.70%	72.20%	74.20%	65.50%	65.10%	63.90%	71.40%	69.80%	60.00%	100.00%	100.00%	100.00%	
Cerebrovascular diseases	33.30%	42.30%	27.10%	33.00%	42.70%	26.50%	34.60%	40.90%	29.60%	29.30%	50.20%	67.20%	65.90%	35.70%	
Chronic lower respiratory diseases	49.20%	53.40%	45.50%	47.70%	52.60%	43.70%	55.40%	56.50%	54.20%	48.50%	50.00%	82.60%	50.00%	57.10%	
Accident (unintentional injury)	78.00%	84.40%	66.70%	76.90%	83.10%	66.20%	82.20%	88.90%	68.90%	70.40%	91.70%	96.00%	96.80%	79.20%	
Motor vehicle accident	94.20%	93.90%	94.90%	93.70%	93.10%	95.20%	95.40%	95.90%	94.30%	91.00%	97.10%	100.00%	99.40%	90.00%	
Falls	34.60%	43.20%	27.60%	34.10%	42.00%	27.70%	37.80%	50.00%	26.90%	30.20%	57.10%	71.40%	75.00%	62.50%	
Accidental poisoning	98.80%	98.90%	98.40%	98.60%	98.80%	98.10%	100.00%	100.00%	100.00%	98.30%	100.00%	100.00%	100.00%	100.00%	
Accidental drowning	92.10%	92.10%	92.30%	90.90%	90.70%	91.30%	100.00%	100.00%	100.00%	84.10%	100.00%	100.00%	100.00%	100.00%	
Influenza and pneumonia	38.40%	46.00%	30.90%	38.70%	46.40%	31.50%	36.70%	44.40%	27.90%	32.40%	66.20%	76.00%	48.90%	42.90%	
Alzheimer's disease	12.60%	18.50%	9.70%	12.60%	17.90%	10.10%	12.20%	22.00%	7.00%	12.00%	16.10%	15.60%	23.50%	30.00%	
Diabetes	59.50%	62.60%	56.00%	58.90%	61.60%	55.80%	61.70%	65.20%	56.90%	52.30%	69.40%	71.90%	82.40%	70.00%	
Intentional self-harm (suicide)	91.00%	90.00%	94.80%	91.20%	90.50%	93.80%	90.50%	88.60%	97.80%	88.60%	98.60%	100.00%	100.00%	90.90%	
Chronic liver disease and cirrhosis	91.50%	93.40%	87.60%	90.90%	93.20%	86.00%	93.30%	93.80%	92.30%	91.10%	88.50%	93.80%	97.10%	75.00%	
Nephritis, nephrotic syndrome and nephrosis	43.60%	43.80%	43.40%	44.80%	47.00%	42.70%	41.00%	37.70%	45.50%	36.30%	54.10%	75.00%	73.30%	33.30%	
Septicemia	57.50%	65.10%	49.80%	58.70%	65.10%	51.90%	53.40%	64.80%	43.50%	51.80%	72.80%	70.00%	75.00%	100.00%	
Assault (homicide)	99.20%	99.30%	98.70%	99.10%	99.20%	98.40%	100.00%	100.00%	100.00%	99.30%	99.60%	96.20%	100.00%	100.00%	
Parkinson's disease	25.80%	28.70%	21.40%	25.00%	27.80%	20.90%	30.90%	34.10%	25.00%	24.50%	30.30%	75.00%	40.00%	50.00%	
Essential (primary) hypertension and hypertensive renal disease	33.30%	41.20%	27.60%	33.40%	41.10%	27.80%	32.70%	41.70%	25.80%	30.70%	40.70%	42.90%	47.40%	0.00%	
Human immunodeficiency virus (HIV) disease	99.30%	99.10%	100.00%	99.20%	99.10%	100.00%	100.00%	100.00%	100.00%	98.80%	100.00%	100.00%	100.00%	100.00%	
INJURY BY FIREARMS	92.10%	91.80%	94.60%	92.90%	92.80%	93.40%	88.80%	87.20%	100.00%	86.50%	99.00%	97.80%	100.00%	100.00%	
DRUG-INDUCED DEATHS	98.40%	98.80%	97.70%	98.20%	98.80%	97.20%	99.10%	98.30%	100.00%	98.00%	100.00%	97.80%	100.00%	100.00%	
ALCOHOL-INDUCED DEATHS	96.30%	96.00%	97.10%	96.20%	95.70%	97.70%	96.80%	97.30%	95.70%	96.00%	95.20%	100.00%	99.10%	80.00%	

* Expected years of life at birth for all U.S. residents.

**Urban=Maricopa, Pima, Pinal and Yuma counties.

Note: The causes of death for 2005 are classified by the Tenth Revision of the International Classification of Diseases (ICD-10).