

**TABLE 3B-4
RATES¹ OF REPORTED CASES OF GONORRHEA, CHLAMYDIA, EARLY SYPHILIS AND GENITAL HERPES
BY AGE AND GENDER, ARIZONA, 2005**

Age Group	GONORRHEA			CHLAMYDIA			EARLY SYPHILIS			GENITAL HERPES		
	Males	Females	Total	Males	Females	Total	Males	Females	Total	Males	Females	Total
0-4	2.2	1.4	1.8	3.5	10.5	6.9	0.0	0.0	0.0	1.3	3.7	2.5
5-9	0.4	0.4	0.4	2.1	3.6	2.8	0.0	0.0	0.0	0.4	0.4	0.4
10-14	4.4	18.5	11.3	16.3	97.1	55.6	0.0	0.0	0.0	0.4	2.3	1.4
15-19	160.8	346.5	250.3	510.1	2619.4	1526.3	3.1	6.2	4.6	13.0	61.5	36.3
20-24	330.2	365.8	347.1	790.4	2913.3	1801.9	10.3	8.8	9.6	34.9	94.8	63.4
25-29	230.4	184.7	208.5	443.0	1223.0	817.2	15.6	8.5	12.2	37.4	59.9	48.2
30-34	150.4	90.1	121.4	227.8	521.4	369.2	15.9	6.7	11.5	29.6	49.6	39.2
35-39	99.7	50.7	75.6	114.6	209.9	161.5	17.0	7.0	12.1	13.2	21.9	17.4
40-44	85.7	32.6	59.4	61.2	102.4	81.6	13.4	5.4	9.4	12.1	24.0	18.0
45-54	39.5	13.2	26.1	32.1	41.2	36.7	8.2	1.6	4.8	8.8	24.6	16.8
55-64	18.4	2.2	9.9	8.8	8.0	8.4	3.2	0.4	1.7	6.4	11.3	9.0
65-over	2.6	1.1	1.8	3.4	2.8	3.1	0.3	0.0	0.1	2.0	4.1	3.2
Total	86.6	77.3	81.9	167.5	535.2	351.8	7.0	3.2	5.1	12.5	26.8	19.7

¹Number of cases per 100,000 population.

Note: The increase in the number of chlamydia and gonorrhea cases reported for 2005 was likely due to changes in surveillance and may not represent a true increase in incidence.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of Infectious Disease Services