

**TABLE 3B-4
RATES¹ OF REPORTED CASES OF GONORRHEA, CHLAMYDIA, EARLY SYPHILIS AND GENITAL HERPES
BY AGE AND GENDER, ARIZONA, 2006**

Age group	GONORRHEA			CHLAMYDIA			EARLY SYPHILIS			GENITAL HERPES		
	Males	Females	Total	Males	Females	Total	Males	Females	Total	Males	Females	Total
0-4	1.2	0.4	0.8	3.7	6.8	5.2	0.0	0.4	0.2	3.7	3.8	3.7
5-9	0.4	2.3	1.3	0.9	4.5	2.7	0.0	0.0	0.0	0.0	0.9	0.4
10-14	3.0	17.9	10.5	10.7	100.3	54.9	0.0	0.0	0.0	0.4	3.1	1.7
15-19	210.2	376.2	291.4	562.8	2895.7	1699.7	4.9	7.5	6.1	14.6	66.9	40.1
20-24	362.7	416.5	389.0	843.0	3147.8	1950.2	18.1	10.7	15.5	40.9	133.9	85.5
25-29	265.3	268.8	267.0	498.6	1409.5	929.8	18.7	9.7	15.3	45.3	101.5	71.9
30-34	167.1	107.4	138.6	220.9	589.2	396.8	17.7	5.7	12.1	36.6	71.6	53.3
35-39	129.1	73.2	102.3	126.8	264.5	193.2	22.7	6.4	14.9	30.5	50.6	40.1
40-44	90.8	38.0	64.8	80.4	125.0	102.4	19.8	3.7	11.9	25.6	35.6	30.6
45-54	50.3	14.8	32.4	28.3	37.1	32.9	10.2	1.0	5.6	14.8	31.8	23.4
55-64	17.5	1.9	9.3	12.3	11.2	11.8	2.1	0.3	1.1	10.3	12.8	11.6
65-over	2.8	0.0	1.3	2.5	3.2	2.9	0.6	0.0	0.3	4.8	2.7	3.6
Total	99.2	91.5	95.3	178.9	593.3	386.1	9.0	3.2	6.2	18.0	37.8	27.9

¹Number of cases per 100,000 population.

Note: Table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2006.

Note: Denominators for unknown or transgender category are not available.

Rates per 100,000 population.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of Infectious Disease Services, STD Program.