

**TABLE 3B-5  
FREQUENCY OF REPORTED CASES, PERCENT DISTRIBUTION AND RATES OF EARLY AND LATE SYPHILIS,  
GONORRHEA, CHLAMYDIA AND HERPES BY ETHNICITY, ARIZONA, 2006**

Race/ethnicity	SYPHILIS						GONORRHEA						CHLAMYDIA			HERPES		
	Early			Late			Resistant			Total			Cases	%	Rate	Cases	%	Rate
	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate						
<b>White Non-Hispanic</b>	134	34.4	3.5	109	20.6	2.9	0	0.0	0.0	1,672	28.1	44.0	5,707	23.7	150.1	687	39.5	18.1
<b>Black or African American</b>	37	9.5	14.9	37	7.0	14.9	0	0.0	0.0	948	15.9	382.4	1,572	6.5	634.2	96	5.5	38.7
<b>Hispanic or Latino</b>	129	33.2	7.6	256	48.3	15.1	0	0.0	0.0	1,384	23.3	81.6	6,900	28.6	406.9	345	19.8	20.3
<b>Asian or Pacific Islander</b>	2	0.5	1.2	5	0.9	3.1	0	0.0	0.0	22	0.4	13.6	155	0.6	95.7	7.0	0.4	4.3
<b>American Indian or Alaska Native</b>	28	7.2	8.4	29	5.5	8.7	0	0.0	0.0	417	7.0	125.7	2,246	9.3	676.8	122	7.0	36.8
<b>Not Specified</b>	59	15.2	N/A	94	17.7	NA	0	0.0	NA	1506	25.3	NA	7,510	31.2	NA	483	27.8	NA
<b>Total</b>	389	100.0	6.2	530	100.0	8.5	0	0.0	0.0	5,949	100.0	95.3	24,090	100.0	386.0	1,740	100.0	27.9

Note: Table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2006.

Note: Rates per 100,000 population.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of Infectious Disease Services, STD Program.