

**TABLE 3B-5  
FREQUENCY OF REPORTED CASES, PERCENT DISTRIBUTION AND RATES OF EARLY AND LATE SYPHILIS,  
GONORRHEA, CHLAMYDIA AND HERPES BY ETHNICITY, ARIZONA, 2007**

Race/ethnicity	SYPHILIS						GONORRHEA						CHLAMYDIA			HERPES		
	Early			Late			Resistant			Total			Cases	%	Rate	Cases	%	Rate
	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate						
<b>White Non-Hispanic</b>	163	28.8	4.2	120	18.5	3.1	0	0.0	0.0	1280	25.3	33.1	5518	22.2	142.5	691	34.5	17.8
<b>Black or African American</b>	43	7.6	17.0	56	8.6	22.1	0	0.0	0.0	876	17.3	345.6	1832	7.4	722.7	130	6.5	51.3
<b>Hispanic or Latino</b>	140	24.8	7.8	278	42.8	15.5	0	0.0	0.0	1087	21.5	60.4	6224	25.0	346.1	390	19.5	21.7
<b>Asian or Pacific Islander</b>	6	1.1	3.5	4	0.6	2.4	0	0.0	0.0	23	0.5	13.5	155	0.6	91.3	18	0.9	10.6
<b>American Indian or Alaska Native</b>	103	18.2	30.5	52	8.0	15.4	0	0.0	0.0	319	6.3	94.4	2167	8.7	641.6	119	5.9	35.2
<b>Not Specified</b>	110	19.5	N/A	140	21.5	N/A	0	0.0	NA	1477	29.2	N/A	8970	36.1	N/A	655	32.7	N/A
<b>Total</b>	565	100.0	8.8	650	100.0	10.1	0	0.0	0.0	5062	100.0	78.7	24866	100.0	386.6	2003	100.0	31.1

Note: Table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2007.

Note: Rates per 100,000 population.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV, HEP C, STD.