

TABLE 3B-4
RATES¹ OF REPORTED CASES OF GONORRHEA, CHLAMYDIA, EARLY SYPHILIS AND GENITAL HERPES
BY AGE AND GENDER, ARIZONA, 2008

Age group	GONORRHEA			CHLAMYDIA			EARLY SYPHILIS			GENITAL HERPES		
	Males	Females	Total	Males	Females	Total	Males	Females	Total	Males	Females	Total
0-4	0.8	1.6	1.2	4.7	9.0	6.8	0.0	0.0	0.0	3.5	5.3	4.4
5-9	0.0	0.4	0.2	2.5	5.3	3.9	0.0	0.0	0.0	1.7	1.8	1.7
10-14	3.0	8.4	5.6	19.0	110.3	63.5	0.0	0.4	0.2	0.0	4.9	2.4
15-19	123.4	214.1	167.3	676.2	3017.7	1809.7	6.0	3.7	4.9	13.3	67.7	39.6
20-24	219.5	256.5	237.0	937.6	3002.1	1914.0	18.0	16.4	17.5	45.7	114.2	78.1
25-29	154.8	110.5	134.1	486.4	1212.0	831.8	29.7	12.7	21.6	51.2	95.2	72.1
30-34	87.1	50.6	69.7	245.6	537.2	386.4	26.8	11.4	19.4	29.3	70.7	49.7
35-39	80.9	38.7	60.4	141.3	269.0	203.3	27.0	6.9	17.2	30.9	64.0	47.2
40-44	44.6	18.0	31.5	79.6	106.5	92.8	29.3	7.2	18.6	24.5	45.4	34.8
45-54	27.3	8.2	17.7	33.1	38.2	35.7	21.3	2.6	11.8	15.3	28.6	22.0
55-64	9.1	0.9	4.9	8.2	9.9	9.1	3.8	0.9	2.3	13.9	16.1	15.2
65-over	2.9	0.9	1.8	3.4	4.1	3.8	0.5	0.0	0.2	4.2	6.2	5.3
Total	57.1	48.3	52.8	195.5	562.8	379.0	13.1	4.4	8.8	18.5	38.2	28.4

¹Number of cases per 100,000 population.

Note: Table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2008.

Note: Denominators for unknown or transgender category are not available.

Rates per 100,000 population.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.