

**TABLE 3B-5
FREQUENCY OF REPORTED CASES, PERCENT DISTRIBUTION AND RATES OF EARLY AND LATE SYPHILIS,
GONORRHEA, CHLAMYDIA AND HERPES BY ETHNICITY, ARIZONA, 2008**

| Race/ethnicity | SYPHILIS | | | | | | GONORRHEA | | | | | | CHLAMYDIA | | | HERPES | | |
|---|----------|-------|------|-------|-------|------|-----------|-----|------|-------|-------|-------|-----------|-------|-------|--------|-------|------|
| | Early | | | Late | | | Resistant | | | Total | | | Cases | % | Rate | Cases | % | Rate |
| | Cases | % | Rate | Cases | % | Rate | Cases | % | Rate | Cases | % | Rate | | | | | | |
| White Non-Hispanic | 229 | 39.8 | 5.8 | 163 | 20.7 | 4.1 | 0 | 0.0 | 0.0 | 827 | 24.0 | 21.0 | 5,158 | 20.8 | 130.9 | 611 | 32.9 | 15.5 |
| Black or African American | 51 | 8.9 | 19.8 | 52 | 6.6 | 20.2 | 0 | 0.0 | 0.0 | 661 | 19.2 | 256.8 | 2,004 | 8.1 | 778.6 | 106 | 5.7 | 41.2 |
| Hispanic or Latino | 202 | 35.1 | 11.1 | 354 | 44.9 | 19.4 | 0 | 0.0 | 0.0 | 903 | 26.2 | 49.5 | 7,537 | 30.4 | 413.3 | 363 | 19.6 | 19.9 |
| Asian or Pacific Islander | 7 | 1.2 | 4.1 | 2 | 0.3 | 1.2 | 0 | 0.0 | 0.0 | 17 | 0.5 | 9.9 | 173 | 0.7 | 100.3 | 7 | 0.4 | 4.1 |
| American Indian or Alaska Native | 55 | 9.6 | 16.1 | 50 | 6.3 | 14.6 | 0 | 0.0 | 0.0 | 251 | 7.3 | 73.3 | 2,533 | 10.2 | 739.8 | 182 | 9.8 | 53.2 |
| Not Specified | 31 | 5.4 | NA | 168 | 21.3 | NA | 0 | 0.0 | NA | 790 | 22.9 | NA | 7,364 | 29.7 | NA | 586 | 31.6 | NA |
| Total | 575 | 100.0 | 8.8 | 789 | 100.0 | 12.1 | 0 | 0.0 | 0.0 | 3,449 | 100.0 | 52.8 | 24,769 | 100.0 | 379.0 | 1,855 | 100.0 | 28.4 |

Note: Table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2008.

Note: Rates per 100,000 population.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.