

# Technical Notes

---

## Sources of Data

## Definitions

- ▶ **URBAN AND RURAL AREAS**
- ▶ **RACE/ETHNICITY**
- ▶ **RATES**
- ▶ **TERMS RELATED TO REPRODUCTIVE HEALTH**
- ▶ **MEDICAL TERMS USED ON THE BIRTH CERTIFICATE**
  - ✓ MEDICAL RISK FACTORS FOR THIS PREGNANCY
  - ✓ COMPLICATIONS OF LABOR AND/OR DELIVERY
  - ✓ ABNORMAL CONDITIONS OF THE NEWBORN
  - ✓ CONGENITAL ANOMALIES OF CHILD
- ▶ **DIAGNOSIS AND PROCEDURE CODES USED TO ANALYZE THE HOSPITAL DISCHARGE DATA**
- ▶ **TERMS RELATED TO MORTALITY**



## SOURCES OF DATA

Information on births, deaths, and fetal deaths is compiled from the original documents filed with the Arizona Department of Health Services', Office of Vital Records and from transcripts of original birth and death certificates filed in other states but affecting Arizona residents. (Copies of certificates for births, deaths and fetal deaths occurring to Arizona residents outside the United States are not sent to Arizona).

Information on induced terminations of pregnancy (abortions) performed on Arizona women is compiled from reports sent to the Arizona Department of Health Services by facilities within but only rarely outside the state.

Pregnancies are the sum of live births, spontaneous terminations of pregnancy (fetal deaths or stillbirths) and induced terminations of pregnancy (abortions).

Arizona has no central registry for marriage and divorce records. Statistics are limited to counts of marriages and divorces reported monthly by the Clerk of the Superior Court in each county in which the marriage or divorce occurred. These reports contain no demographic data on the people marrying or divorcing.

Data on morbidity, levels of disease and disability in the population, are obtained for certain infectious diseases that must be reported by law. The ADHS Epidemiology and Disease Control Services conduct surveillance and monitoring of these reportable diseases and provided data for the morbidity sections in this report.

All short-stay nonfederal hospitals in Arizona are required to submit every six months uniform patient reports to the Arizona Department of Health Services. The Section of Cost Reporting and Discharge Data Review in the Bureau of Public Health Statistics collects the information about both hospital inpatient discharges and emergency room visits.

Population denominators for Arizona residents, used to calculate rates for 1980, 1990, and 2000, are census enumerations from the U.S. Census Bureau. Population denominators for the intercensal years utilize the best available data sources. They are described in detail at <http://www.azdhs.gov/plan/menu/info/pd.htm>

Our website at <http://www.azdhs.gov/plan/index.htm> provides additional reports and studies, as well as links to other sources of information on topics such as morbidity from certain diseases or population composition.

## DEFINITIONS

### Urban and Rural Areas

For the purpose of this report, the following are Arizona's **urban areas**: Phoenix-Scottsdale-Mesa Metropolitan Statistical Area (Maricopa and Pinal Counties), Tucson Metropolitan Statistical Area (Pima County), and Yuma Metropolitan Statistical Area (Yuma County). The remaining counties (Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Mohave, Navajo, Santa Cruz, and Yavapai) comprise Arizona's **rural areas**. The 1994 list of Metropolitan Statistical Areas and their component counties for the United States is available at:

<http://www.cdc.gov/nchs/data/dvs/mcd/geog94msa.txt>

### Race/Ethnicity

Race/ethnic designations used in this report are White non-Hispanic, Hispanic or Latino, Black or African-American, American Indian or Alaska Native (including Aleut and Eskimo) and Asian or Pacific Islander (including Hawaiian). In some of the trend tables the designation "Other" includes Asian and Pacific Islanders. Non-White Hispanics are included in their appropriate race groups.

The American Indian figures include those living both on and off the reservation.

The race/ethnicity of mother, father or decedent is as stated on the certificate.

Implementation of the revised certificates of live birth, death and fetal death in Arizona is necessary in order to provide compatibility with the new federal standards for collecting and presenting data on race and Hispanic origin as established by the Office of Management and Budget (OMB) in October 1997. These federal standards require that when self-identification is used, a method for reporting more than one race should be adopted. In addition, the Asian or Pacific Islander category will be split into two categories – one called "Asian" and the other called "Native Hawaiian or Other Pacific Islander".

### Rates

Rate is a measure of the frequency of some event in relation to a unit of population during a specified time period such as a year; events in the numerator of the year occur to individuals in the denominator. Rates express the likelihood (or risk) of the event in the specified population during a particular time and are generally expressed as units of population in the denominator (per 1,000, 10,000, 100,000 and so forth).

Many rates shown in this report are based on a small population, a small number of events or both. Rates based on small numbers are unreliable and thus should be viewed with caution. Rates for many counties or Arizona's ethnic minorities also vary considerably from year to year due to small populations and few events.

### Terms Related to Reproductive Health

**Abortion Rate** - Number of reported abortions to females of all ages during a calendar year per 1,000 females of childbearing age (15-44 years).

**Abortion Ratio** - Number of abortions reported during a period per 1,000 live births occurring during the same period.

**Birth or Live Birth** - The complete expulsion or extraction from its mother of a product of human conception, irrespective of the duration of pregnancy which, after such expulsion or extraction, breathes or shows any other evidence of life such as beating of the heart, pulsation of the umbilical cord or definite movement of voluntary muscles, whether or not the umbilical cord has been cut or the placenta is attached. Heartbeats are to be distinguished from transient cardiac contractions; respirations are to be distinguished from fleeting respiratory efforts or gasps.

**Birth rate** - Number of live births during a calendar year per 1,000 population.

**Birth weight** - The weight of a neonate determined immediately after delivery or as soon thereafter as possible.

**Fetal death** - Death prior to the complete expulsion or extraction from its mother of a product of human conception, irrespective of the duration of pregnancy and which is not an induced termination of pregnancy. The death is indicated by the fact that, after such expulsion or extraction, the fetus does not breathe or show any other evidence of life, such as beating of the heart, pulsation of the umbilical cord or definite movement of voluntary muscles. Heartbeats are to be distinguished from transient cardiac contractions; respirations are to be distinguished from fleeting respiratory efforts or gasps.

**Fetal death rate** - Number of fetal deaths at 20 or more weeks of gestation per 1,000 live births + fetal deaths.

**Fertility rate (general)** - Total number of live births to women of all ages during a calendar year per 1,000 women of childbearing age (15-44 years old).

**Fertility rate (total)** - The sum of age-specific birth rates of women at each age group 10-14 through 45-49. Since five-year age groups are used, the sum is multiplied by five.

**Infant death** - Any death at any time from birth up to, but not including, the first year of age (364 days, 23 hours, 59 minutes from the moment of birth).

**Infant mortality rate** - Number of infant deaths per 1,000 live births.

**Low-birthweight** - Any neonate weighing less than 2,500 grams at birth (less than 5 pounds 8 ounces).

**Maternal mortality rate** - Number of deaths attributed to maternal conditions (i.e. related to or aggravated by pregnancy or its management) per 100,000 live births.

**Neonatal death** - Death of a live born neonate before the neonate becomes 28 days old (up to and including 27 days, 23 hours, 59 minutes from the moment of birth).

**Neonatal mortality rate** - Number of neonatal deaths per 1,000 live births.

**Perinatal mortality rate** - The sum of infant deaths of less than 7 days and fetal deaths with a stated or presumed period of gestation of 28 weeks or more per 1,000 live births + fetal deaths of 28 or more weeks of gestation..

**Postneonatal death** - Any death of a live born infant at least 28 days of age but less than one year of age.

**Postneonatal mortality rate** - Number of postneonatal deaths per 1,000 live births.

**Post term** - Any neonate whose birth occurs from the beginning of the first day (295<sup>th</sup> day) of the 43<sup>rd</sup> week following onset of the last menstrual period.

**Pregnancy rate** - The sum of live births, fetal deaths and induced terminations of pregnancy per 1,000 women of childbearing age (15-44 years old).

**Preterm** - Any neonate whose birth occurs through the end of the last day of the 37<sup>th</sup> week (259<sup>th</sup> day), following onset of the last menstrual period.

**Term** - Any neonate whose birth occurs from the beginning of the first day (260<sup>th</sup> day) of the 38<sup>th</sup> week, through the end of the last of the 42<sup>nd</sup> week (294<sup>th</sup> day), following onset of the last menstrual period.

**Very Low-Birthweight** - Any neonate whose weight at birth is 1,500 grams or less (less than 3 pounds 5 ounces).

## Medical Terms Used on the Birth Certificate

### Medical risk factors for this pregnancy

**Anemia** - Hemoglobin level of less than 10.0 g/dL during pregnancy, or hematocrit of less than 30 percent during pregnancy.

**Cardiac disease** - Disease of the heart.

**Acute or chronic lung disease** - Disease of the lungs during pregnancy.

**Diabetes** - Metabolic disorder characterized by excessive discharge of urine and persistent thirst; includes juvenile onset, adult onset and gestational diabetes during pregnancy.

**Genital herpes** - Infection of the skin of the genital area by herpes simplex virus.

**Hydramnios/Oligohydramnios** - Any noticeable excess (hydramnios) or lack (oligohydramnios) of amniotic fluid.

**Hemoglobinopathy** - A blood disorder caused by alteration in the genetically determined molecular structure of hemoglobin (for example, sickle cell anemia).

**Hypertension, chronic** - Blood pressure persistently greater than 140/90, diagnosed prior to onset of pregnancy or before the 20th week of gestation.

**Hypertension, pregnancy-associated** - An increase in blood pressure of at least 30 mm hg systolic or 15 mm hg diastolic on two measurements taken 6 hours apart after the 20th week of gestation.

**Eclampsia** - The occurrence of convulsions and/or coma unrelated to other cerebral conditions in women with signs and symptoms of preeclampsia.

**Incompetent cervix** - Characterized by painless dilation of the cervix in the second trimester or early in the third trimester of pregnancy, with premature expulsion of membranes through cervix and ballooning of the membranes into vagina, followed by rupture of the membrane and subsequent expulsion of the fetus.

**Previous infant 4,000+ grams** - The birthweight of a previous live-born child was over 4,000 grams (8lbs.14oz.).

**Previous preterm or small-for-gestational-age-infant** - Previous birth of an infant prior to term (before 37 completed weeks of gestation) or of an infant weighing less than the 10th percentile for gestational age using a standard weight for age chart.

**Renal disease** - Kidney disease.

**Rh Sensitization** - The process or state of becoming sensitized to the Rh factor as when an Rh-negative woman is pregnant with an Rh-positive fetus.

**Uterine bleeding** - Any clinically significant bleeding during the pregnancy taking into consideration the stage of pregnancy; any second or third trimester bleeding or third trimester bleeding of the uterus prior to the onset of labor.

### Complications of labor and/or delivery

**Febrile** - A fever greater than 100 degrees F. or 38 degrees C. occurring during labor and/or delivery.

**Meconium, moderate/heavy** - Meconium consists of undigested debris from swallowed amniotic fluid, various products of secretion and shedding by the gastrointestinal tract, moderate to heavy amounts of meconium in the amniotic fluid noted during labor and/or delivery.

**Premature rupture of membranes (more than 12 hours)** - Rupture of membranes at any time during pregnancy and more than 12 hours before the onset of labor.

**Abruptio placenta** - Premature separation of normally implanted placenta from the uterus.

**Placenta previa** - Implantation of the placenta over or near the internal opening of the cervix.

**Other excessive bleeding** - The loss of significant amount of blood from conditions other than abruptio placenta or placenta previa.

**Seizures during labor** - Maternal seizures occurring during labor from any cause.

**Precipitous labor** (less than 3 hours) Extremely rapid labor and delivery lasting less than 3 hours.

**Prolonged labor** (more than 20 hours) - Abnormally slow progress of labor lasting more than 20 hours.

**Dysfunctional labor** - Failure to progress in a normal pattern of labor.

**Breech/Malpresentation** - At birth, the presentation of the fetal buttocks, rather than the head or other malpresentation.

**Cephalopelvic disproportion** - The relationship of the size, presentation and position of the fetal head to the maternal pelvis, which prevents dilation of the cervix and/or descent of the fetal head.

**Cord prolapse** - Premature expulsion of the umbilical cord in labor before the fetus is delivered.

**Anesthetic complications** - Any complication during labor and/or delivery brought on by an anesthetic agent or agents.

**Fetal distress** - Signs indicating fetal hypoxia (deficiency in amount of oxygen reaching fetal tissues).

#### **Abnormal conditions of the newborn**

**Anemia** - Hemoglobin level of less than 13.0 g/dL, or a hematocrit of less than 39 percent.

**Birth injury** - Impairment of the infant's body function or structure due to adverse influences that occurred at birth.

**Fetal alcohol syndrome** - A syndrome of altered prenatal growth and development occurring in infants born of women who consumed excessive amounts of alcohol during pregnancy.

**Hyaline membrane disease/RDS** - A disorder primarily of prematurity, manifested clinically by respiratory distress and pathologically by pulmonary hyaline membranes and incomplete expansion of the lungs at birth.

**Meconium aspiration syndrome** - Aspiration of meconium by the fetus or newborn, affecting the lower respiratory system.

**Assisted ventilation (less than 30 minutes)** - A mechanical method of assisting respiration for newborns with respiratory failure.

**Assisted ventilation (30 minutes or more)** - Newborn placed on assisted ventilation for 30 minutes or longer.

**Seizures** - A seizure of any etiology.

#### **Congenital anomalies of child**

**Anencephalus** - Absence of the cerebral hemispheres.

**Spina bifida/meningocele** - Developmental anomaly characterized by defective closure of the bony encasement of the spinal cord, through which the cord and meninges may or may not protrude.

**Hydrocephalus** - Excessive accumulation of cerebrospinal fluid within the ventricles of the brain with consequent enlargement of cranium.

**Microcephalus** - A significantly small head.

**Other central nervous system anomalies** - Other specified anomalies of the brain, spinal cord, and nervous system.

**Heart malformations** - Congenital anomalies of heart.

**Other circulatory/respiratory anomalies** - Other specified anomalies of the circulatory and respiratory systems.

**Rectal atresia/stenosis** - Congenital absence, closure, or narrowing of the rectum.

**Tracheo-esophageal fistula/Esophageal atresia** - An abnormal passage between the trachea and the esophagus; esophageal atresia is the congenital absence or closure of the esophagus.

**Omphalocele/Gastroschisis** - An omphalocele is a protrusion of variable amounts of abdominal viscera from a midline defect at the base of the umbilicus. In gastroschisis, the abdominal viscera protrude through an abdominal wall defect, usually on the right side of the umbilical cord insertion.

**Other gastrointestinal anomalies** - Other specified congenital anomalies of the gastrointestinal system.

**Malformed genitalia** - Congenital anomalies of the reproductive organs.

**Renal agenesis** - One or both kidneys are completely absent.

**Other urogenital anomalies** - Other specified congenital anomalies of the organs concerned in the production and excretion of the urine, together with organs of reproduction.

**Cleft lip/palate** - Cleft lip is a fissure or elongated opening of the lip; cleft palate is a fissure in the roof of the mouth. These are failures of embryonic development.

**Polydactyly/Syndactyly/Adactyly** - Polydactyly is the presence of more than five digits on hands and/or feet; syndactyly is having fused or webbed fingers and/or toes; adactyly is the absence of fingers and/or toes.

**Club foot** - Deformities of the foot, which is twisted out of shape or position.

**Diaphragmatic hernia** - Herniation of the abdominal contents through the diaphragm into the thoracic cavity usually resulting in respiratory distress.

**Down's syndrome** - The most common chromosomal defect with most cases resulting from an extra chromosome (trisomy 21).

## Diagnosis and Procedure Codes Used to Analyze the Hospital Discharge Data

Diagnostic groupings and code numbers used in Chapter 4 and Chapter 7 are based on the International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM).

The tabulations of the hospital inpatient data by first-listed diagnosis and all-listed procedures utilize the diagnostic categories available respectively at <http://www.azdhs.gov/plan/hip/cat/icd9primary.xls> and <http://www.azdhs.gov/plan/hip/cat/icd9procedure.xls>.

The ICD-9-CM diagnostic categories used to identify specific mental disorders are available online at <http://www.azdhs.gov/plan/hip/for/mental/2007/mental107.xls>

### Terms Related to Mortality

The most common, and perhaps the most valuable measure of the likelihood (or risk) of death in the specified population during a particular time is the **crude death rate**. It is computed as the number of deaths per 1,000 or 100,000 population.

**Age-adjusted mortality rates** - Because mortality from most causes of death occurs predominately among the elderly, a population group with a larger proportion of older persons would have a higher mortality rate. The "age-adjustment" removes the effect of the age differences among sub-populations (or in the same population over time) by placing them all in a population with a standard age distribution. All age-adjusted mortality rates in this report were computed by the direct method that is by weighting the age-specific rates for a given year by the age distribution of a standard population. The weighted age-specific rates are then added to produce the summary rate for all ages combined. Beginning with the 2000 data year, a new population standard for the age adjustment of mortality rates has replaced the standard based on the 1940 population and used since 1943. The new standard uses the age composition of the 2000 U.S. projected population. The standard is expressed in terms of a "standard million": the relative distribution of the 2000 population of the United States totaling 1 million in 10-year age groups:

Age group	2008 population	2008 deaths	Age-specific rates in 2008	2000 standard	Age-adjusted rate for 2008
A	B	C	D (C/B)*100000	E	F D*E
<1	98995	625	631.3	.013818	8.7
1-4	402486	128	31.8	.055317	1.8
5-14	927978	134	14.4	.145565	2.1
15-24	903796	754	83.4	.138646	11.6
25-34	949725	900	94.8	.135573	12.8
35-44	898425	1472	163.8	.162613	26.6
45-54	844161	3403	403.1	.134834	54.4
55-64	659678	5496	833.1	.087247	72.7
65-74	432451	7663	1772.0	.066037	117.0
75-84	307069	11947	3890.7	.044842	174.5
85+	110157	12604	11441.9	.015508	177.4
All ages	6534921	45128		TOTAL	<b>659.6</b>

The age-adjusted mortality rates should be viewed as relative indexes rather than as actual measures of mortality risk. It is also important to note that Age-adjusted rates can only be compared to other age-adjusted rates that use the same population standard.

In this report, all age-adjusted mortality rates are based on the 2000 standard, and they CANNOT BE compared to rates using the 1940 standard population.

**Age-specific mortality** - Number of deaths in a specific age group during a calendar year.

**Alcohol-induced deaths** - This category was expanded in 2003. Causes of death attributable to alcohol mortality include mental and behavioral disorders due to alcohol use, degeneration of nervous system due to alcohol use, alcoholic polyneuropathy, alcoholic cardiomyopathy, alcoholic gastritis, alcoholic liver disease, finding of alcohol in blood, accidental poisoning by and exposure to alcohol, intentional self-poisoning by alcohol, poisoning by alcohol, undetermined intent.

**Cause of death** - For the purpose of national mortality statistics, every death is attributed to one underlying condition, based on information reported on the death certificate and utilizing the international rules for selecting the underlying cause of death from the reported conditions.

**Cause-specific mortality** - Number of deaths from a specified cause during a calendar year.

**Classification of causes of death** - The cause of death used in this report is the underlying cause classified according to the *International Classification of Diseases (ICD)*. Beginning with the 2000 data year in Arizona (1999 nationally), a new revision of the International Classification of Diseases was implemented. The Tenth Revision (ICD-10) has replaced the Ninth Revision (ICD-9), which was in effect since 1979.

**Comparability ratios** - Comparability ratios are measures of comparison between ICD-9 and ICD-10. Any comparison of causes of mortality in Arizona before and after the implementation of ICD-10 needs to take into account the changes in statistical trends that can be attributed to changes in the classification system alone. Comparability ratio of 1.0 indicates that the same number of deaths would be assigned to a cause-of-death when ICD-9 or ICD-10 was used. Comparability-modified number of deaths and mortality rates are shown for the four causes of death for which the discontinuity in trend is substantial (influenza and pneumonia, Alzheimer's disease, nephritis, or septicemia). The following online table provides comparable cause-of-death categories in ICD-9 and ICD-10 and shows comparability ratios for 113 selected causes of death:

<http://www.azdhs.gov/plan/report/ahs/ahs2007/pdf/compratio.pdf>

**Drug-induced deaths** - This category was expanded in 2003. Causes of death attributable to drug-related mortality include mental and behavioral disorders due to psychoactive substance use, accidental poisoning by and exposure to drugs, suicide by drugs, homicide by drugs and poisoning by drugs, undetermined intent

**Firearm mortality** - Causes of death attributable to firearm mortality include accidental discharge of firearms, suicide by firearms, homicide by firearms, legal intervention involving discharge of firearms, terrorism involving firearms and discharge of firearms, undetermined intent.



# INDEX TO TABLES

(Numbers and letters refer to tables)

## A

- Abdominal pain**, emergency room visits for, 4C-2, 7C-2
- Abnormal conditions of the newborn** 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-34, 5B-28, 5B-30
- Abortion**  
Abortions per 1,000 live births 1D-1, 1D-2, 5D-2  
Abortions per 1,000 women 1A-1, 1A-2, 1A-3, 1A-4, 1A-5, 1A-6, 1A-7, 5A-1  
Age group 1A-2, 1A-3, 1A-4, 1A-5, 1A-6, 1A-7, 1D-1, 1D-3, 1D-5, 5A-4, 5D-4  
County of residence 5A-1, 5A-4, 5D-1, 5D-2, 5D-3, 5D-4, 5D-5, 5D-6, 5D-7  
Gestation, length of 1D-1, 1D-3, 1D-4, 5D-7  
Hospitalized for complications 1D-1  
Marital status 5D-6  
Number performed in Arizona 1D-1, 5D-3  
Race/ethnicity 1A-1, 1A-2, 1A-3, 1A-4, 1A-5, 1A-6, 1D-1, 1D-2, 5D-5  
Reason for termination 1D-1  
Type of facility 1D-1
- Abruptio placenta**, (*see also* **Complications of labor and/or delivery**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-27
- Abstinence from cigarette smoking** 6A-1, 6B-2-1
- Accidents**, (*see* **Injuries, unintentional**)
- Acute myocardial infarction**, deaths from 2B-6, 2C-28, 2C-29, 5E-12
- Adverse effects of medical treatment**, emergency room visits related to, 4D-2, 7D-2
- Adverse effects of medical treatment**, inpatient discharges related to, 4D-1, 7D-1
- African-American residents**, (*see* **Black residents**)
- Age at death**  
County of residence 5E-11  
Gender differences 2D-1, 2D-3,  
Race/ethnic differences 2D-1, 2D-3,  
For the leading causes 2D-3  
Urban/rural differences 2D-3
- AIDS**, (*see* **HIV/AIDS**)
- Alcohol-induced deaths** 2B-6, 2C-28, 2C-29, 5E-12, 9C
- Alcoholic liver disease** 2B-6, 2C-28, 2C-29, 5E-12
- Alcoholic psychoses**, emergency room visits for, 4C-1, 7C-1
- Alcoholic psychoses**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1
- Alcohol dependence syndrome**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1
- Alcohol use during pregnancy** 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25, 5B-30
- Alzheimer's disease** deaths from 2B-1, 2B-2, 2B-4, 2B-5, 2B-6, 2C-27, 2C-28, 2D-3, 2D-4, 5E-11, 5E-12, 5E-13, 5E-14, 5E-33, 5E-34
- American Indian or Alaska Native residents** 1A-1, 1A-2, 1A-3, 1A-4, 1A-5, 1A-6, 1B-2, 1B-3, 1B-4, 1B-5, 1B-12, 1B-13, 1B-20, 1B-22, 1B-23, 1B-24, 1B-25, 1B-26, 1B-28, 1B-30, 1B-31, 1B-32, 1B-33, 1B-34, 1B-35, 1C-4, 1C-5, 1D-1, 1D-2, 1D-4, 1D-5, 2A-1, 2B-3, 2B-4, 2C-4, 2C-11, 2C-15, 2C-19, 2C-23, 2C-27, 2D-1, 2D-2, 2D-3, 2D-4, 3B-5, 3C-3, 5B-7, 5B-8, 5B-10, 5B-15, 5B-18, 5B-22, 5C-5, 5D-5, 5E-6, 5E-18
- Amebiasis**, 3A-1, 3A-2, 5F-1, 5F-2
- Amniocentesis**, (*see also* **Obstetric procedures**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-26
- Anemia**  
Mother's condition 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25  
Newborn's condition 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-28  
Deaths from 2B-6, 2C-28, 5E-12
- Anencephalus**, (*see also* **Congenital anomalies**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 1B-35, 5B-29
- Anesthetic complications** 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-27
- Anxiety states**, emergency room visits for, 4C-1, 7C-1
- Anxiety states**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1
- Aortic aneurysm and dissection**, 2B-6, 2C-28, 2C-29, 5E-12
- Appendix, diseases of**, deaths from 2B-6, 2C-28, 2C-29, 5E-12
- Appendicitis**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1
- Appendectomy**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1
- Appendix, diseases of** (deaths from) 2B-6, 2C-28, 2C-29, 5E-12
- Arizona Health Care Cost Containment System (AHCCCS)** 1B-25, 1B-26, 1B-27, 1B-28, 1B-29, 1B-32, 1B-33, 1B-34, 5B-20
- Arterio- & angiocardiology using contrast material**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1
- Artificial rupture of membranes**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1
- Asian (Asian or Pacific Islander) residents** 1B-14, 1B-15, 1B-20, 1B-22, 1B-23, 1B-24, 1B-26, 1B-29, 1B-30, 1B-31, 1B-32, 1B-33, 1B-34, 1B-37, 1D-1, 1D-2, 1D-4, 1D-5, 2A-1, 2B-3, 2B-4, 2C-4, 2C-11, 2C-15, 2C-19, 2C-23, 2C-27, 2D-3, 2D-4, 3B-5, 3C-3, 5B-7, 5B-8, 5B-10, 5B-15, 5B-18, 5D-5, 5E-6, 5E-18
- Assault**, (*see* **Homicide**)
- Assisted ventilation**, (*see also* **Abnormal conditions of the newborn**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-28
- Asthma**, deaths from 2B-6, 2C-28, 2C-29, 6A-8, 5E-12
- Asthma**, emergency room visits for, 4C-1, 4C-2, 7C-1, 7C-2
- Asthma**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1
- Atelectasis** deaths from 2C-7, 5E-20
- Atherosclerosis**, deaths from 2B-6, 2C-28, 2C-29, 5E-12
- Atherosclerosis, coronary**, inpatient discharges with 4A-1, 4A-2, 4A-3, 4A-5, 7A-1
- Atherosclerotic cardiovascular disease**, deaths from 2B-6, 2C-28, 5E-12
- Attendant at birth** 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15
- Axial tomography, computerized**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

## B

- Bilateral destruction/occlusion of fallopian tubes**, inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1
- Births**  
Abnormal conditions of the newborn 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-34, 5B-28, 5B-30  
Age of father 1B-19  
Age of mother 1A-2, 1A-3, 1A-4, 1A-5, 1A-6, 1B-1, 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15, 1B-19,



Brain 2B-6, 2B-8, 2C-28, 5E-11  
Breast 2B-2, 2B-3, 2B-4, 2B-6, 2B-8, 2C-28, 6A-5, 5E-11, 5E-12, 5E-13, 5E-14, 6B-5  
Cervix uteri 2B-4, 2B-5, 2B-6, 2B-8, 5E-11, 6A-5, 6B-5  
Colon, rectum, and anus (colorectal) 2B-4, 2B-5, 2B-6, 2B-8, 2C-28, 5E-11, 6A-5, 6B-5  
Corpus uteri and uterus 2B-6, 2B-8, 2C-28, 5E-11  
Esophagus 2B-6, 2B-8, 2C-28, 5E-11  
Hodgkin's disease 2B-6, 2B-8, 2C-28, 5E-11  
In situ 2B-6, 2C-28, 5E12  
Kidney 2B-6, 2B-8, 2C-28, 5E-11  
Larynx 2B-6, 2C-28, 5E-11  
Leukemia 2B-6, 2C-28, 5E-11  
Lip, oral cavity and pharynx 2B-6, 2C-28, 5E-11  
Liver and intrahepatic bile ducts 2B-6, 2C-28, 5E-11  
Lung 2B-2, 2B-3, 2B-4, 2B-6, 2B-8, 2C-28, 6A-5, 5E-11, 5E-12, 5E-13, 5E-14, 6B-5  
Lymphoid and hematopoietic tissue 2B-6, 2C-28, 5E-11  
Malignant lymphoma other than Hodgkin's disease 2B-8  
Melanoma of skin 2B-8, 6A-5  
Multiple myeloma 2B-8  
Oropharyngeal 2B-6, 2B-8, 2C-28, 5E-11  
Ovary 2B-6, 2C-28, 5E-11  
Pancreas 2B-6, 2B-8, 2C-28, 5E-11  
Prostate 2B-4, 2B-12, 2B-7, 5E-11, 5E-12, 5E-13, 5E-14, 6A-5, 6B-5  
Skin 2B-4, 2B-5, 2B-6, 2B-8, 2C-28, 5E-11, 5E-12  
Stomach 2B-6, 2B-8, 2C-28, 5E-11  
Trachea, bronchus and lung 2B-6, 2B-8, 2C-28, 5E-11

**Cancer** (malignant neoplasm), inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1  
Breast 4A-1, 4A-2, 4A-3, 4A-5, 7A-1  
Large intestine 4A-1, 4A-2, 4A-3, 4A-5, 7A-1  
Prostate 4A-1, 4A-2, 4A-3, 4A-5, 7A-1  
Trachea, bronchus and lung 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Cancer** (neoplasm), emergency room visits for, 4C-1, 7C-1

**Cardiac catheterization**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

**Cardiac disease** as a medical risk factor for pregnancy 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25

**Cardiac dysrhythmias**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Cardiovascular disease**, deaths from 2B-2, 2B-3, 2B-4, 2B-5, 2B-6, 2C-28, 2D-3, 2D-4, 6A-5-32, 5E-11, 5E-12, 5E-13, 5E-14, 6B-5-29

**Cellulitis and abscess**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Cephalopelvic disproportion**, (*see also* **Complications of labor/delivery**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-27

**Cerebrovascular disease** (stroke), deaths from 2B-1, 2B-2, 2B-3, 2B-4, 2B-5, 2B-6, 2C-28, 2D-3, 2D-4, 6A-5-32, 5E-11, 5E-12, 5E-13, 5E-14, 6B-5-28

**Cerebrovascular disease**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Certain conditions originating in the perinatal period**, deaths due to 2B-6, 2C-28, 5E-11

**Cesarean deliveries**, (*see also* **Delivery, method of**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-26, 5B-30

**Cesarean section**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

**Chest pain**, emergency room visits for, 4C-2, 7C-2

**Chickenpox** 3A-1, 3A-2, 5F-1, 5F-2

**Chinese mothers**, number of births to, 1B-22

**Chlamydia** 3B-1, 3B-3, 3B-4, 3B-5, 6A-2, 5F-1, 5F-2

**Cholecystectomy**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

**Cholelithiasis**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Cholelithiasis and other disorders of gallbladder**, deaths from 2B-6, 2C-28, 5E-11

**Cholera** 3A-1, 5F-1, 5F-2

**Chronic liver disease and cirrhosis** deaths from 2B-1, 2B-3, 2B-4, 2B-5, 2B-6, 2C-28, 5E-11, 5E-12, 5E-13, 5E-14

**Chronic lower respiratory diseases** deaths from 2B-1, 2B-2, 2B-3, 2B-4, 2B-5, 2B-6, 2C-28, 2D-3, 2D-4, 5E-11, 5E-12, 5E-13, 5E-14

**Chronic obstructive pulmonary diseases**, (*see* **Chronic lower respiratory diseases**)

**Cigarette smoking during pregnancy** 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25, 5B-30

**Circulatory system, diseases of**, emergency room visits for, 4C-1, 7C-1

**Circulatory system, diseases of**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Cirrhosis**, deaths from 6A-10, 6B-10

**Cleft lip/palate**, (*see also* **Congenital anomalies**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 1B-35, 5B-29

**Club foot**, (*see also* **Congenital anomalies**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 1B-35, 5B-29

**Coccidioidomycosis (Valley Fever)**  
Reported cases 3A-1, 5F-1, 5F-2  
Deaths from 3A-2

**Cold, exposure to**, deaths from 2B-9

**Colorado Tick Fever** 3A-1

**Complications of medical and surgical care**, deaths due to 2B-6, 2C-28, 5E-12

**Complications of medical and surgical care**, visits related to 4D-2, 7D-2

**Complications of medical and surgical care**, inpatient discharges related to, 4D-1, 7D-1

**Communicable diseases**, (*see* **Diseases, reportable**)

**Complications of labor and/or delivery**  
1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15, 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-34, 5B-27, 5B-30

**Complications of pregnancy, childbirth and the puerperium**, deaths from 1C-2, 2B-6, 2C-28, 5E-11

**Complications, surgical and medical**, emergency room visits for, 4C-1, 7C-1

**Conditions originating in the perinatal period**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Congenital anomalies** (birth defects) **reported on birth certificate** 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 1B-35, 5B-29

**Congenital malformations, deformations and chromosomal abnormalities** deaths from 2B-6, 2C-28, 5E-12

**Congenital anomalies**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Contusion with intact skin surface**, emergency room visits for, 4C-2, 7C-2

**Contusions**, emergency room visits for, 4C-1, 7C-1

**Cord prolapse**, (*see also* **Complications of labor/delivery**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-27

**Coronary artery bypass graft**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

**Coronary artery obstruction, removal of**, (inpatient operation), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

**Coronary heart disease**, deaths from 2B-4, 2B-5, 5E-11, 5E-13, 5E-14, 6A-7, 6B-7

**Cuban mothers**, number of births to, 1B-22

**Cystoscopy with or without biopsy**, (inpatient operation), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

## D

**Day of birth** 1B-18

**Day of death** 2A-3

**Deaths** 2A-1, 2A-2, 2A-3, 2A-4, 2B-1, 2B-2, 2B-3, 2B-4, 2-5, 2B-6, 2B-7, 2B-8, 2B-9, 2B-10, 2B-11, 1C-1, 2C-2, 2C-3, 2C-4, 2C-5, 2C-6, 2C-7, 2C-8, 2C-9, 2C-10, 2C-11, 2C-12, 2C-13, 2C-14, 2C-15, 2C-16, 2C-17, 2C-18, 2C-19, 2C-20, 2C-21, 2C-22, 2C-23, 2C-24, 2C-25, 2C-26, 2C-27, 2C-28, 2D-1, 2D-2, 2D-3, 2D-4, 6A-3, 6A-4, 6A-5, 5E-1, 5E-2, 5E-3, 5E-4, 5E-5, 5E-6, 5E-7, 5E-8, 5E-9, 5E-10, 5E-11, 5E-12, 5E-13, 5E-14, 5E-15, 5E-16, 5E-17, 5E-18, 5E-19, 5E-20, 5E-21, 5E-22, 5E-23, 5E-24, 5E-25, 5E-26, 5E-27, 5E-28, 5E-29, 5E-30, 5E-31, 5E-32, 5E-33, 5E-34, 6B-3, 6B-4, 6B-5, 9B, 9C

Age adjusted rates 2B-2, 2B-3, 2B-4, 2B-5, 5E-11

Age-specific numbers and rates 2C-5, 2C-6, 2C-7, 2C-8, 2C-9, 2C-10, 2C-11, 2C-12, 2C-13, 2C-14, 2C-15, 2C-16, 2C-17, 2C-18, 2C-19, 2C-20, 2C-21, 2C-22, 2C-23, 2C-24, 2C-25, 2C-26, 2C-27, 2C-28, 2C-29, 5E-20, 5E-25, 5E-26, 5E-27, 5E-28, 5E-29, 5E-30, 5E-31, 5E-32, 5E-33, 5E-34, 9B

Average/median age at death 2D-1, 2D-3, 5E-11

County of residence 5E-11, 5E-12, 5E-13, 5E-14

Crude death rates 2B-6, 5E-13

Gender differences 2B-2, 2B-3, 2B-4, 2B-5, 2B-6, 2D-3, 2D-4

Premature mortality 2D-2, 2D-4

Race/ethnicity 2B-3, 2B-4, 2D-3, 2D-4

Urban/rural differences 2B-5, 2D-3, 2D-4

**Decedent status** 2A-4, 2B-10, 5E-8

**Delivery, method of** 1B-26, 1B-28, 1B-32, 1B-33, 1B-34

**Dengue**, 3A-1, 3A-2, 5F-1, 5F-2

**Depression**, emergency room visits for, 4C-1, 7C-1

**Depression**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Diabetes mellitus** as medical risk factor for pregnancy, 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25, 5B-26

**Diabetes mellitus**, deaths from 2B-1, 2B-2, 2B-3, 2B-4, 2B-5, 2B-6, 2C-28, 2C-29, 2D-3, 2D-4, 5E-11, 5E-12, 5E-13, 5E-14

**Diabetes mellitus**, emergency room visits for, 4C-1, 7C-1

**Diabetes mellitus**, inpatient discharges with 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Diabetes-related deaths** 6A-6, 6B-6

**Diaphragmatic hernia**, (*see also* **Congenital anomalies**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 1B-35, 5B-29

**Digestive system, diseases of**, emergency room visits for, 4C-1, 7C-1

**Digestive system, diseases of**, inpatient discharges with 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Diseases, reportable**

AIDS 3C-1, 3C-2, 3C-3, 3C-4, 6A-2, 5F-3

Amebiasis 3A-1, 5F-1, 5F-2,

Botulism 3A-1, 5F-1, 5F-2

Brucellosis 3A-1, 3A-2, 5F-1, 5F-2

Central nervous system 3A-1, 5F-1, 5F-2

Chickenpox 3A-1, 5F-1

Chlamydia 3B-1, 3B-3, 3B-4, 3B-5, 5F-1, 5F-2

Coccidioidomycosis (Valley Fever) 3A-1, 5F-1, 5F-2

Colorado Tick Fever 3A-1, 5F-1, 5F-2

Cryptosporidiosis 3A-1, 5F-1, 5F-2

Dengue 3A-1, 3A-2, 5F-1, 5F-2

E. coli O157:H7 3A-1, 5F-1, 5F-2

Encephalitis, viral 3A-1, 5F-1, 5F-2

Enteric 3A-1, 5F-1, 5F-2

Erllichiosis 3A-1

Fungal 3A-1, 5F-1, 5F-2

Giardiasis 3A-1, 5F-1, 5F-2

Gonorrhea 3B-1, 3B-3, 3B-4, 3B-5, 6A-2, 5F-1, 5F-2, 6B-2

Gonococcal Pelvic Inflammatory Disease 3B-1, 5F-1, 5F-2

Haemophilus influenzae b 3A-1, 6A-2-16, 5F-1, 5F-2

Hantavirus Pulmonary Syndrome 3A-1, 5F-1, 5F-2

Hepatitis 3A-1, 6A-2-16, 5F-1, 5F-2

Legionellosis 3A-1, 3A-2, 5F-1, 5F-2

Listeriosis 3A-1, 3A-2, 5F-1, 5F-2

Lyme disease 3A-1, 5F-1, 5F-2

Malaria 3A-1, 3A-2, 5F-1, 5F-2

Measles 3A-1, 6A-2-15, 5F-1, 5F-2

Meningitis, aseptic 3A-1, 5F-1, 5F-2

Meningitis, bacterial 3A-1, 5F-1, 5F-2

Meningitis, meningococcal 3A-1, 5F-1, 5F-2

Methicillin Resistant S. aureus (invasive) 3A-1, 5F-1, 5F-2

Mumps 3A-1, 6A-2-15, 5F-1, 5F-2

Plague 3A-1, 5F-1, 5F-2

Rabies, Human 3A-1, 5F-1, 5F-2

Reyes Syndrome 3A-1, 5F-1, 5F-2

Rocky Mountain Spotted Fever 3A-1, 3A-2, 5F-1, 5F-2

Rubella 3A-1, 5F-1, 5F-2

Salmonella paratyphi A 3A-1, 5F-1, 5F-2

Salmonella paratyphi B 3A-1, 5F-1, 5F-2

Salmonella paratyphi C 3A-1, 5F-1, 5F-2

Salmonellosis (except S. typhi and S. paratyphi) 3A-1, 5F-1, 5F-2

Sexually transmitted 3B-1, 5F-1, 5F-2

Shigellosis 3A-1, 5F-1, 5F-2

St. Louis encephalitis 3A-1, 5F-1, 5F-2

Streptococcal-Group A 3A-1, 5F-1, 5F-2

Streptococcal-Group B 3A-1, 5F-1, 5F-2

Streptococcus pneumoniae 3A-1, 5F-1, 5F-2

Syphilis 3B-1, 3B-3, 3B-4, 3B-5, 6A-2-14, 5F-1, 5F-2

Congenital 6A-2-14

Toxic Shock Syndrome 3A-1, 5F-1, 5F-2

Tuberculosis 3A-1, 6A-2-16, 5F-1, 5F-2

Pulmonary Tuberculosis 3A-1, 5F-1, 5F-2

Tularemia 3A-1, 5F-1, 5F-2

Typhoid fever 3A-1, 5F-1, 5F-2

Vaccine preventable 3A-1, 5F-1, 5F-2

Vibrio spp. 3A-1, 5F-1, 5F-2

Vancomycin, resistant Enterococcus spp. 3A-1,

West Nile virus 3A-1, 5F-1, 5F-2

Yersiniosis 3A-1, 5F-1, 5F-2

Zoonotic 3A-1, 5F-1, 5F-2

**Disorders relating to short gestation and unspecified low-birthweight**, (*see also* **Infant mortality**) 2C-5, 2C-6, 2C-7, 5E-20

**Dissolutions of marriage** 5G-4, 5G-5, 5G-6

**Diverticula of intestine**

**Divorces**, (*see* **Dissolutions of marriage**)

**Downs syndrome**, (*see also* **Congenital anomalies**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 1B-35, 5B-29

**Drowning and submersion, accidental**, deaths from 2B-4, 2B-5, 2B-6, 2C-8, 2C-9, 2C-11, 2C-15, 2C-28, 2C-29, 5E-11, 5E-12, 5E-13, 5E-14  
Among children 4 years and younger 6A-3-23

**Drug abuse, nondependent**, inpatient discharges for, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Drug abuse, nondependent**, emergency room visits for, 4C-1, 7C-1

**Drug dependence**, emergency room visits for, 4C-1, 7C-1

**Drug dependence**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Drug-induced deaths** 2B-6, 2C-28, 2C-29, 5E-12, 9C

**Drug psychoses**, emergency room visits for, 4C-1, 7C-1  
**Drug psychoses**, inpatient discharges with 4A-1, 4A-2, 4A-3, 4A-5, 7A-1  
**Dysfunctional labor**, (*see also Complications of labor/delivery*) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-27

## E

**E. coli O157:H7** 3A-1, 5F-1, 5F-2  
**Eclampsia**, (*see also Medical risk factors for pregnancy*) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25  
**Electronic fetal monitoring**, (*see also Obstetric procedures*) 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-34, 5B-26, 5B-30  
**Emphysema**, deaths from 2B-6, 2C-28, 5E-12  
**Encephalitis, arthropod-borne**, deaths from 2B-6, 2C-28, 5E-12  
**Encephalitis, viral** 3A-1, 3A-2, 5F-1, 5F-2  
**Endocarditis, acute and subacute**, deaths from 2B-6, 2C-28, 5E-12  
**Endocrine nutritional metabolic and immunity diseases**, emergency room visits for, 4C-1, 7C-1  
**Endocrine nutritional metabolic and immunity diseases**, inpatient discharges with 4A-1, 4A-2, 4A-3, 4A-5, 7A-1  
**Endoscopy of small intestine**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1  
**Endoscopy of large intestine**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1  
**Endotracheal tube, insertion of** (inpatient procedure),  
**Enteritis and colitis**, noninfectious, inpatient discharges with 4A-1, 4A-2, 4A-3, 4A-5, 7A-1  
**Enterocolitis due to *Clostridium difficile*** inpatient discharges with 4A-1, 4A-2, 4A-3, 4A-5, 7A-1  
**Enterococcus spp., Vancomycin resistant**, 3A-1  
**Episiotomy**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1  
**Erlinchiosis** 3A-1, 5F-1, 5F-2  
**Events of undetermined intent** as cause of death 2B-6, 2C-28, 5E-12  
**Excision (partial) of bone**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1  
**Excision (or destruction) of intervertebral disc**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1  
**Excision (partial) of large intestine**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

## F

**Falls**, deaths from 2B-4, 2B-5, 2B-6, 2B-9, 2C-23, 2C-27, 2D-3, 2D-4, 6A-4, 5E-11, 5E-12, 5E-13, 5E-14, 5E-31, 5E-32, 5E-33, 5E-34, 6B-4  
**Father's age** 1B-19  
**Father's education** 1B-21  
**Father's race/ethnicity** 1B-20  
**Febrile**, (*see also Complications of labor/delivery*) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-27  
**Fertility**, (*see also Births*) 1A-1, 1A-2, 1A-3, 1A-4, 1A-5, 1A-6  
    General fertility 1B-1  
    Total fertility 1B-1  
**Fetal alcohol syndrome** 1B-26, 1B-27, B-28, 1B-32, 1B-33, 1B-34, 5B-28, 5B-30  
**Fetal deaths**, (*see also Perinatal mortality*) 1C-3, 1C-4, 1C-5, 5A-4, 5C-1, 5C-2, 5C-3, 5C-4, 5C-5, 5C-6  
**Fever**, emergency room visits for, 4C-2, 7C-2  
**Fetal distress**, (*see Complications of labor/delivery*) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-27

**Filipino mothers**, number of births to, 1B-22  
**Fire, flames, smoke (exposure to)**, deaths due to 2B-9, 2B-6, 2C-28, 5E-12, 6A-3  
    Among children 4 years and younger 6A-3  
    Among persons 65 years and over 6A-3  
**Firearms, accidental discharge of**, deaths due to 2B-6, 2B-9, 2C-28, 5E-12  
**Firearms, injury by**, deaths due to 2B-6, 2C-28, 6A-4, 5E-12, 6B-4  
**Foreign bodies**, emergency room visits for, 4C-1, 7C-1  
**Fractures, all sites**, emergency room visits for, 4C-1, 4C-2, 7C-1, 7C-2  
**Fractures, excluding lower limb**, emergency room visits for, 4C-2, 7C-2  
**Fracture, reduction of**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1  
**Fractures**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1  
**Fungal diseases** 3A-1, 3A-2, 5F-1, 5F-2

## G

**Gastric band (gastric restrictive procedure)**, 4B-1, 4B-2, 4B-3, 4B-4, 7B-1  
**Gastroenteritis and colitis**, emergency room visits for, 4C-2, 7C-2  
**Genitourinary system, diseases of**, emergency department visits for, 4C-1, 7C-1  
**Genitourinary system, diseases of**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1  
**Gestation, length of**  
    At birth 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15, 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-22, 5B-23, 5B-24, 5B-30  
    At termination of pregnancy 1D-1, 1D-3, 1D-4, 5D-7  
**Giardiasis** 3A-1, 3A-2, 5F-1, 5F-2  
**Glomerulonephritis, chronic**, deaths from 2B-6, 2C-28, 2C-29, 5E-12  
**Gonococcal Pelvic Inflammatory Disease**, 3B-1, 3B-2, 5F-1, 5F-2  
**Gonorrhea** 3B-1, 3B-2, 3B-3, 3B-4, 3B-5, 6A-2, 5F-1, 5F-2, 6B-2  
**Gross reproduction rate** 1B-1

## H

**Haemophilus influenzae (invasive)** 3A-1, 6A-2, 5F-1, 5F-2, 6B-2  
**Hanta Pulmonary Syndrome** 3A-1, 3A-2, 5F-1, 5F-2  
**Hawaiian mothers**, number of births to, 1B-22  
**Headache**, emergency room visits for, 4C-2, 7C-2  
**Health Objectives for the Year 2010** 6A-1, 6A-2, 6A-3, 6A-4, 6A-5, 6A-4, 6A-5, 6A-6, 6A-7, 6A-8, 6A-9, 6A-10, 6B-1, 6B-2, 6B-3, 6B-4, 6B-5, 6B-6, 6B-7, 6B-8, 6B-9, 6B-10  
    Cancer 6A-5, 6B-5  
    Diabetes 6A-6, 6B-6  
    Heart disease and stroke 6A-7, 6B-7  
    Human immunodeficiency virus (HIV) disease 6A-9, 6B-9  
    Injury and violence 6A-4, 6B-4  
    Maternal, infant and child health 6A-1, 6B-1  
    Respiratory diseases 6A-8, 6B-8  
    Responsible sexual behavior 6A-2, 6B-2  
    Substance abuse 6A-10, 6B-10  
    Vaccine preventable diseases 6A-3, 6B-3  
**Heart disease**, deaths from 2B-1, 2B-2, 2B-3, 2B-4, 2B-5, 2B-6, 2B-7, 2C-28, 2D-3, 2D-4, 5E-11, 5E-12,

5E-13, 5E-14

**Heart disease, excluding ischemic**, emergency room visits for, 4C-2, 7C-2

**Heart disease**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, &A-1

**Heart failure**, death from, 2B-6, 2C-28, 5E-12

**Heart failure, congestive**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Heart malformations**, (*see also* **Congenital anomalies**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 1B-35, 5B-29

**Heat, exposure to**, deaths from 2B-9

**Hemodialysis**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

**Hemoglobinopathy**, (*see also* **Medical risk factors for pregnancy**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25

**Hepatitis** 3A-1, 3A-2, 6A-2, 5F-1, 5F-2, 6B-2

**Hernia**, deaths from 2B-6, 2C-28, 5E-12

**Herpes, genital** (reported cases) 3B-1, 5F-1, 5F-2  
As a medical risk factor for pregnancy 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25

**Hip replacement**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

**Hispanic (Latino) residents** 1A-1, 1A-2, 1A-3, 1A-4, 1A-5, 1A-6, 1B-2, 1B-3, 1B-4, 1B-5, 1B-8, 1B-9, 1B-20, 1B-22, 1B-23, 1B-24, 1B-25, 1B-26, 1B-28, 1B-30, 1B-31, 1B-32, 1B-33, 1B-34, 1B-35, 1C-4, 1C-5, 1D-1, 1D-2, 1D-4, 1D-5, 2A-1, 2B-3, 2B-4, 2C-4, 2C-11, 2C-15, 2C-19, 2C-23, 2C-27, 2D-1, 2D-2, 2D-3, 2D-4, 3B-5, 3C-3, 5B-7, 5B-8, 5B-10, 5B-15, 5B-18, 5B-22, 5C-5, 5D-5, 5E-6, 5E-18

**HIV Disease/AIDS**  
Reported cases 3C-1, 3C-2, 3C-3, 3C-4, 6A-9, 5F-3, 6B-9  
Deaths from 2B-4, 2B-5, 2B-6, 2C-16, 2C-17, 2C-19, 2C-28, 6A-9, 5E-11, 5E-12, 5E-13, 5E-14, 5E-29, 5E-30, 6B-9

**Hodgkin's disease**, deaths from 2B-6, 2C-28, 5E-12

**Homicide (assault)** as cause of death, 2B-1, 2B-2, 2B-3, 2B-4, 2B-5, 2B-6, 2B-11, 2C-7, 2C-8, 2C-9, 2C-11, 2C-12, 2C-13, 2C-15, 2C-16, 2C-17, 2C-19, 2C-23, 2C-28, 6A-4, 5E-11, 5E-12, 5E-13, 5E-14, 5E-25, 5E-26, 5E-27, 5E-28, 5E-29, 5E-30, 5E-31, 5E-32, 6B-4, 9C

**Homicide (assault)**, emergency room visits related to, 4D-2, 7D-2

**Homicide (assault)**, inpatient discharges related to, 4D-1, 7D-1

**Hyaline membrane disease/Respiratory Distress Syndrome**, (*see also* **Abnormal conditions of the newborn**) 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-34, 5B-28, 5B-30

**Hydramnios/Oligohydramnios**, (*see also* **Medical risk factors for pregnancy**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25

**Hydrocephalus**, (*see also* **Congenital anomalies**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 1B-35, 5B-29

**Hyperplasia of prostate** 2B-6, 2C-28, 5E-12

**Hypertension, chronic**, (*see also* **Medical risk factors for pregnancy**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25

**Hypertension, essential (primary), and Hypertensive renal disease** deaths from 2B-6, 2C-28, 5E-12

**Hypertension, pregnancy-associated**, (*see also* **Medical risk factors for pregnancy**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25

**Hysterectomy**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

**Indian Health Service (IHS)** 1B-26, 1B-27, 1B-28, 1B-29, 1B-32, 1B-33, 1B-34, 5B-22

**Induced terminations of pregnancy**, (*see* **Abortions**)

**Induction of labor**, (*see also* **Obstetric procedures**) 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-34, 5B-26

**Infant mortality**, (*see also* **Perinatal mortality**) 2C-1, 2C-2, 2C-3, 2C-4, 2C-5, 2C-6, 2C-7, 2C-28, 6A-1, 5E-15, 5E-16, 5E-17, 5E-18, 5E-19, 5E-20, 5E-21, 5E-22, 5E-23, 5E-24, 6B-2-1, 8C-1, 8C-2

**Infection of urinary tract**, emergency room visits for, 4C-2, 7C-2

**Infections of kidney**, deaths from 2B-6, 2C-28, 5E-12

**Infectious and parasitic diseases**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Infectious and parasitic diseases**, emergency room visits for, 4C-1, 7C-1

**Inflammatory diseases of female pelvic organs**, deaths from 2B-6, 2C-28, 5E-12

**Influenza**, deaths from 2B-6, 2C-28, 5E-12

**Influenza and pneumonia**, deaths from 2B-1, 2B-2, 2B-4, 2B-5, 2B-6, 2C-5, 2C-6, 2C-7, 2C-11, 2C-19, 2C-23, 2C-27, 2C-28, 2C-29, 2D-3, 2D-4, 5E-11, 5E-12, 5E-13, 5E-14, 5E-20, 5E-25, 5E-26, 5E-29, 5E-30, 5E-31, 5E-32, 5E-33, 5E-34

**Injuries, intracranial**, emergency room visits for, 4C-1, 7C-1

**Injuries of undetermined intent**, emergency room visits, 4D-2, 7D-2

**Injuries of undetermined intent**, inpatient discharges with, 4D-1, 7D-1

**Injuries, superficial**, emergency room visits for, 4C-1, 4C-2, 7C-1, 7C-2

**Injuries, unintentional (accidents)** deaths from 2B-1, 2B-2, 2B-3, 2B-4, 2B-5, 2B-6, 2B-9, 2B-10, 2B-11, 2C-5, 2C-6, 2C-7, 2C-8, 2C-9, 2C-11, 2C-12, 2C-13, 2C-15, 2C-16, 2C-17, 2C-19, 2C-20, 2C-21, 2C-23, 2C-27, 2C-28, 2C-29, 6A-3, 5E-11, 5E-12, 5E-13, 5E-14, 5E-20, 5E-25, 5E-26, 5E-27, 5E-28, 5E-29, 5E-30, 5E-31, 5E-32, 5E-33, 5E-34, 6B-5, 9C  
Accidents involving aircraft 2B-9  
Adverse effects of drugs in therapeutic use 2B-9  
Bee (hornet, wasp) sting 2B-9  
Bitten by a dog 2B-9  
Choked on food 2B-9  
Choked on other object 2B-9  
Cold, exposure to 2B-9  
Dog bite 2B-9  
Drowning 2B-4, 2B-5, 2B-6, 2B-9, 2C-8, 2C-9, 2C-11, 2C-28, 2D-3, 2D-4, 6A-3-23, 5E-11, 5E-12, 5E-13, 5E-14, 5E-25,, 5E-26, 5E-27, 5E-28  
Electric current 2B-9  
Falls 2B-4, 2B-5, 2B-6, 2B-9, 2C-23, 2C-27, 2D-3, 2D-4, 6A-4, 5E-11, 5E-12, 5E-13, 5E-14, 5E-31, 5E-32, 5E-33, 5E-34, 6B-4  
Fire, flames, smoke 2B-9  
Firearm discharge 2B-6, 2B-9, 2C-28, 2C-29, 5E-12  
Heat, exposure to 2B-9  
Inhalation or ingestion of food 2B-9  
Inhalation or ingestion of other objects 2B-9  
Lightning 2B-9  
Misadventures to patients during surgical and medical care 2B-9  
Motorcyclist 2B-10  
Motor vehicle accidents 2B-1, 2B-2, 2B-3, 2B-4, 2B-5, 2B-6, 2B-9, 2B-9, 2B-10, 2C-8, 2C-9, 2C-11, 2C-12, 2C-13, 2C-15, 2C-16, 2C-17, 2C-19, 2C-20, 2C-21, 2C-23, 2C-27, 2C-28, 2D-3, 2D-4, 6A-3-18, 5E-1, 5E-12, 5E-13, 5E-14, 5E-25, 5E-26, 5E-27, 5E-28, 5E-29, 5E-30, 5E-31, 5E-32, 5E-33, 5E-34, 6B-3-18  
Overexertion 2B-9  
Poisoning by drugs 2B-9

**Incompetent cervix**, (*see also* **Medical risk factors for pregnancy**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25

Poisoning by other substances, gases and vapors 2B-9  
Railway accidents 2B-9  
Spider bite 2B-9  
Storms and floods 2B-9  
Suffocation and other threats to breathing 2B-9  
Venomous snakes and lizards 2B-9  
Water transport accidents 2B-9

**Injuries, unintentional (accidents),**

emergency room visits for, 4D-2, 7D-2  
Cutting or piercing instruments 4D-2, 7D-2  
Drowning/submersion, 4D-2, 7D-2  
Falls, 4D-2, 7D-2  
Fire and flames, 4D-2, 7D-2  
Firearm missile, 4D-2, 7D-2  
Foreign body entering eye or other orifice, 4D-2, 7D-2  
Machinery, 4D-2, 7D-2  
Motor vehicle accident, 4D-2, 7D-2  
Natural and environmental factors, 4D-2, 7D-2  
Overexertion, 4D-2, 7D-2  
Pedal cycle, 4D-2, 7D-2  
Poisoning by drugs, 4D-2, 7D-2  
Suffocation, 4D-2, 7D-2

**Injuries, unintentional (accidents), inpatient discharges with,** 4D-1, 7D-1

Falls, 4D-1, 7D-1  
Fire and flames, 4D-1, 7D-1  
Machinery, 4D-1, 7D-1  
Motor vehicle accident, 4D-1, 7D-1  
Natural and environmental factors, 4D-1, 7D-1  
Poisoning by drugs, 4D-1, 7D-1

**Injury and poisoning, emergency room visits for,** 4C-1, 4D-2, 7C-1, 7D-2

**Inpatient deaths** 2A-4, 2B-11, 5E-8

**Intentional self-harm, see Suicide**

**Intrauterine hypoxia and birth asphyxia** 2C-5, 2C-6, 2C-7, 5E-20

**Invertebral disc disorders, inpatient discharges for,** 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Ischemic heart disease, deaths from** 2B-6, 2C-28, 5E-12

**J**

**Japanese mothers, number of births to,** 1B-22

**K**

**Knee replacement, (inpatient procedure),** 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

**L**

**Leading causes of death, (see also, Deaths),** 2B-1, 2B-2, 2B-3, 2B-4, 2B-5, 2B-6, 2C-28, 5E-11, 5E-12, 5E-13, 5E-14

Age adjusted rates 2B-2, 2B-3, 2B-4, 2B-5, 5E-11  
Average/median age at death 2D-1, 2D-3, 5E-11  
County of residence 5E-11, 5E-12, 5E-13, 5E-14  
Crude death rates 2B-6, 5E-13  
Gender differences 2B-2, 2B-3, 2B-4, 2B-5, 2B-6, 2D-3, 2D-4  
Premature mortality 2D-2, 2D-4  
Race/ethnicity 2B-3, 2B-4, 2D-3, 2D-4  
Urban/rural differences 2B-5, 2D-3, 2D-4

**Leading causes of death among**

**infants <1 year** 2C-5, 2C-6, 2C-7, 5E-20  
**children 1-14 years** 2C-8, 2C-9, 2C-10, 2C-11, 5E-25, 5E-26

**adolescents 15-19 years** 2C-12, 2C-13, 2C-14, 2C-15, 5E-27, 5E-28

**young adults 20-44 years** 2C-16, 2C-17, 2C-18, 2C-19, 5E-29, 5E-30

**middle-aged adults 45-64** 2C-20, 2C-21, 2C-22, 2C-23, 5E-31, 5E-32

**elderly 65 years and older** 2C-24, 2C-25, 2C-26, 2C-27, 2C-28, 5E-33, 5E-34

**Selected 113 causes** 2B-6, 2C-28, 5E-12

**Legal intervention, deaths due to** 2B-6, 2C-28, 5E-12

**Legionellosis** 3A-1, 3A-2, 5F-1, 5F-2

**Length of stay (average) for inpatient discharges,** 4A-5

**Leukemia, deaths from** 2B-6, 2C-28, 5E-12

**Listeriosis** 3A-1, 3A-2, 5F-1, 5F-2

**Live births, (see Births)**

**Liver disease and cirrhosis, chronic, deaths from** 2B-3, 2B-4, 2B-5, 2B-6, 2C-28, 5E-11, 5E-12, 5E-13, 5E-14

**Low-birthweight (LBW) births** 1B-2, 1B-3, B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15, 1B-16, 1B-25, 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-34, 5B-16, 6A-1-7, 5B-17, 5B-18, 5B-19, 5B-20, 5B-21, 5B-22, 5B-23, 5B-24, 6B-1,, 9A

**Lung disease, chronic or acute, (see also Medical risk factors for pregnancy)** 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25

**Lyme disease** 3A-1, 5F-1, 5F-2

**Lysis of peritoneal adhesions, (inpatient procedure),** 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

**M**

**Malaria, deaths from,** 2B-6, 2C-28, 2C-29, 5E-12

**Malaria, reported cases,** 3A-1, 3A-2, 5F-1, 5F-2

**Malformed genitalia, (see also Congenital anomalies)** 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 1B-35, 5B-29

**Malignant neoplasm, (see Cancer)**

**Malnutrition, deaths from** 2B-6, 2C-28, 5E-12

**Manic-depressive disorders, emergency room visits for,** 4C-1, 7C-1

**Manic-depressive disorders, inpatient discharges with,** 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Marital status of mother** 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15, 1B-23, 1B-25, 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-24, 5B-14, 5B-15

**Marriages** 5G-1, 5G-2, 5G-3

**Maternal deaths** 1C-1, 1C-2

**Measles**

Number of reported cases 3A-1, 6A-2, 5F-1, 5F-2, 6B-2

Number of deaths 2B-6, 2C-28, 3A-2, 5E-12

**Meconium aspiration syndrome, (see also**

**Abnormal conditions of the newborn)** 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-34, 5B-28

**Meconium, moderate/heavy, (see also**

**Complications of labor/delivery)** 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-27

**Medical risk factors for pregnancy** 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15, 1B-25, 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25, 5B-30

**Meningitis** 2B-6, 2C-28, 3A-1, 3A-2, 5E-12, 5F-1, 5F-2

**Meningococcal infection** 2B-6, 2C-28, 3A-1, 3A-2, 6A-2, 5E-12, 5F-1, 5F-2, 6B-2

**Mental disorders, emergency room visits for,** 4C-1, 4C-2, 7C-1, 7C-2

**Mental disorders, inpatient discharges with,** 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Methicillin Resistant S. aureus (invasive)** 3A-1, 5F-1, 5F-2

**Mexican mothers**, number of births to, 1B-22

**Microcephalus**, (see also **Congenital anomalies**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 1B-35, 5B-29

**Migraine**, emergency room visits for, 4C-2, 7C-2

**Misadventures to patients during surgical or medical care**, death due to 2B-9

**Miscarriages**, (see **Fetal Deaths**)

**Month of birth** 1B-18, 1B-17, 5B-6

**Month of death** 2A-2, 2A-3, 5E-7

**Mortality**, (see **Deaths**)

**Mother's age** 1A-2, 1A-3, 1A-4, 1A-5, 1A-6, 1B-1, 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15, 1B-19, 1B-23, 1B-24, 1B-29, 1B-30, 1B-31, 1B-32, 1B-33, 1B-34, 5A-4, 5B-9, 5B-10

**Mother's education** 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15, 1B-26, 1B-28, 1B-32, 5B-13, 5B-15

**Mother's race/ethnicity** 1A-1, 1A-2, 1A-3, 1A-4, 1A-5, 1A-6, 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15, 1B-20, 1B-22, 1B-23, 1B-24, 1B-25, 1B-26, 1B-29, 1B-30, 1B-31, 1B-32, 1B-33, 1B-34, 5B-8, 5B-10, 5B-15

**Mother's weight gain during pregnancy** 1B-26, 1B-28, 1B-32, 1B-33, 1B-34

**Multiple births** 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15, 1B-16

**Multiple myeloma and immunoproliferative neoplasms**, deaths from 2B-6, 2C-28, 5E-12

**Mumps** 3A-1, 3A-2, 6A-2, 5F-1, 5F-2, 6B-2

**Musculoskeletal system, diseases of**, emergency room visits for, 4C-1, 7C-1

**Musculoskeletal system, diseases of**, inpatient discharges with 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Myocardial infarction, acute**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

## N

**Nausea, vomiting**, emergency room visits for, 4C-2, 7C-2

**Nativity**, (see **Births**)

**Native American**, (see **American Indian**)

**Natural increase from births** 5B-1

**Neonatal deaths** 2C-1, 2C-3, 2C-4, 2C-7, 5C-1, 5E-23, 5E-24

**Nephritis, nephrotic syndrome, and nephrosis**, deaths from 2B-1, 2B-2, 2B-4, 2B-5, 2C-28, 2D-3, 2D-4, 5E-11, 5E-12, 5E-13, 5E-14

**Nervous system, diseases of**, emergency department visits for, 4C-1, 7C-1

**Nervous system, diseases of**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-

**Neuroses (neurotic disorders)**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Newborn's health**

Abnormal conditions 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-34, 5B-28, 5B-30

Admitted to intensive care units 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 5B-22, 5B-24

Congenital anomalies 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 1B-35, 5B-29

Low birth weight 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13,

1B-14, 1B-15, 1B-16, 1B-25, 1B-26, 1B-27, 1B-32, 1B-33, 1B-34, 5B-16, 5B-17, 5B-18,

5B-19, 5B-20, 5B-21, 5B-22, 5B-23, 5B-24, 5B-30

Prematurity 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15, 1B-25, 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-34, 5B-22, 5B-23, 5B-24, 5B-30, 9A

Very low birth weight 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15, 1B-26, 1B-28, 1B-32

**Non-Hodgkin's lymphoma**, deaths from 2B-6, 2C-28, 2C-29, 5E-12

**Nontransport accidents**, deaths from 2B-6, 2C-28, 2C-29, 5E-12

**Notifiable diseases**, (see **Diseases reportable**)

**Nutritional deficiencies**, deaths from 2B-6, 2C-28, 2C-29, 5E-12

## O

**Objectives**, (see **Health objectives for the Year 2010**)

**Obstetric laceration, repair of** (inpatient operation), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

**Obstetric procedures** 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-34, 5B-26, 5B-30

**Omphalocele**, (see also **Congenital anomalies**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 1B-35,

**Oophorectomy and salpingo-oophorectomy**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

**Open wound, excluding head**, emergency room visits for, 4C-2, 7C-2

**Open wound of head**, emergency room visits for, 4C-2, 7C-2

**Operations of war**, deaths from 2B-6, 2C-28, 5E-12

**Operations (inpatient procedures)** on the cardiovascular system 4B-1, 4B-2, 4B-3, 4B-4, 7B-1 digestive system 4B-1, 4B-2, 4B-3, 4B-4, 7B-1 ear 4B-1, 4B-2, 4B-3, 4B-4, 7B-1 endocrine system 4B-1, 4B-2, 4B-3, 4B-4, 7B-1 eye 4B-1, 4B-2, 4B-3, 4B-4, 7B-1 female genital organs 4B-1, 4B-2, 4B-3, 4B-4, 7B-1 hemic and lymphatic system 4B-1, 4B-2, 4B-3, 4B-4, 7B-1 integumentary system 4B-1, 4B-2, 4B-3, 4B-4, 7B-1 male genital organs 4B-1, 4B-2, 4B-3, 4B-4, 7B-1 musculoskeletal system 4B-1, 4B-2, 4B-3, 4B-4, 7B-1 nervous system 4B-1, 4B-2, 4B-3, 4B-4, 7B-1 nose, mouth, pharynx 4B-1, 4B-2, 4B-3, 4B-4, 7B-1 respiratory system 4B-1, 4B-2, 4B-3, 4B-4, 7B-1 urinary system 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

**Otitis media and eustachian tube disorders**, emergency room visits for, 4C-2, 7C-2

**Osteoarthritis and allied disorders**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Outpatient deaths** 2A-4, 2B-11, 5E-8

## P

**Pacific Islanders**, (see **Asian residents**)

**Parkinson's disease**, deaths from 2B-6, 2C-27, 2C-28, 5E-12, 5E-33, 5E-34

**Payee for birth** 1B-26, 1B-27, 1B-28, 1B-29, 1B-32, 1B-33, 1B-34, 6B-7, 5B-20

**Pedal cyclist deaths** 2B-10

**Pedestrian deaths** 2B-10

**Peptic ulcer**, deaths from 2B-6, 2C-28, 5E-12

**Pericardium and (acute) myocarditis**, deaths from, 2B-6, 2C-28, 5E-12

**Perinatal deaths** 1G-3, 6A-1, 5C-1, 6B-2-1

**Pertussis (whooping cough)**

Reported cases, 3A-1, 5F-1, 5F-2

Deaths from 2B-6, 2C-28, 2C-29, 3A-2, 5E-12

**Pharyngitis, acute**, emergency room visits for, 4C-2, 7C-2

**Placenta previa**, (*see also* **Complications of labor/delivery**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-27

**Place of birth/delivery** 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15

**Place of death** 2A-4, 2B-10, 5E-8

**Plague** 3A-1, 3A-2, 5F-1, 5F-2

**Plurality** (*see* **Multiple births**)

**Pneumoconioses and chemical effects** deaths from 2B-6, 2C-28, 5E-12

**Pneumonia** deaths from 2B-6, 2C-28, 5E-12

**Pneumonia**, emergency room visits for, 4C-1, 7C-1

**Pneumonia**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Pneumonitis due to solids and liquids**, deaths from 2B-6, 2C-28, 5E-12

**Poisoning, accidental**, death from, 2B-6, 2B-9, 6A-4, 6B-4

**Poisoning**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Poisonings**, emergency room visits for, 4C-1, 7C-1

**Poliomyelitis, acute**, deaths from 2B-6, 2C-28, 2C-29, 5E-12

**Polydactyly/Syndactyly/Adactyly**, (*see also*

**Congenital anomalies**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 1B-35, 5B-29

**Population denominators** 10A-1, 10A-2, 10B-1, 10B-2, 10C-1

**Postneonatal deaths** 2C-1, 2C-3, 2C-4, 2C-7, 5E-21, 5E-22

**Precipitous labor**, (*see also* **Complications of labor/delivery**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-27

#### **Pregnancies**

Among girls 14 and younger 1A-7, 5A-4

Among girls 17 and younger 1A-2, 5A-3

Among girls 15-17 1A-3, 5A-4

Among girls 15-19 1A-7, 5A-3

Among girls 18-19 1A-4, 5A-4

Among girls 19 and younger 1A-5, 1A-7, 5A-2

By county of residence 5A-1, 5A-2, 5A-3, 5A-4

By outcome 1A-1, 1A-2, 1A-3, 1A-4, 1A-5, 1A-6, 1A-7, 5A-1, 5A-4

By race/ethnicity 1A-1, 1A-2, 1A-3, 1A-4, 1A-5, 1A-6

By women's age 1A-6, 1A-7, 5A-4

**Pregnancy, childbirth and the puerperium**, deaths related to 1C-2, 2B-6, 2C-28, 5E-12

**Pregnancy with abortive outcome**, deaths from 2B-6, 2C-28, 5E-12

**Premature mortality** 2D-2, 2D-4

#### **Prenatal care**

Average number of visits 5B-22

Number of visits 1B-2, 1B-3, 1B-4, 1B-5, 1B-6,

1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13,

1B-14, 1B-15, 1B-25, 1B-26, 1B-27, 1B-28,

1B-31, 1B-32, 1B-33, 1B-34, 5B-12, 5B-21, 9A

Trimester of entry 1B-1, 1B-2, 1B-3, 1B-4, 1B-5,

1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12,

1B-13, 1B-14, 1B-15, 1B-25, 1B-26, 1B-27,

1B-28, 1B-30, 1B-32, 1B-33, 1B-34, 5B-11, 9A

**Preterm births** 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15, 1B-25, 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-34, 5B-22, 6A-1, 5B-23, 5B-24, 5B-30, 6B-2-1

**Previous infant 4,000+ grams** (*see also*

**Medical risk factors for pregnancy**) 1B-26,

1B-28, 1B-32, 1B-33, 1B-34, 5B-25

**Previous preterm or small for gestational age (SGA) infant** (*see also* **Medical risk factors for pregnancy**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25

**Prolonged labor**, (*see also* **Complications of labor/delivery**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-27

**Prostate cancer**, deaths from 2B-2, 2B-3, 2B-4, 2B-5, 2B-6, 2C-27, 2C-28, 2D-3, 2D-4, 5E-11, 5E-12, 5E-13, 5E-14, 5E-33, 5E-34

**Prostatectomy**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

**Psychoses**, inpatient discharges with 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Puerto-Rican mothers**, number of births to, 1B-22

## **R**

**Rabies** 3A-1, 3A-2, 5F-1, 5F-2

#### **Rates**

Age-adjusted death rates 2B-2, 2B-3, 2B-4, 2B-5, 5E-5, 5E-11

Age-specific death rates 2C-1, 2C-2, 2C-3, 2C-6, 2C-7, 2C-8, 2C-9, 2C-10, 2C-12, 2C-13, 2C-14, 2C-16, 2C-17, 2C-18, 2C-20, 2C-21, 2C-22, 2C-24, 2C-25, 2C-26, 52C-29, E-26, 5E-28, 5E-30, 5E-32, 5E-34

Birth rates 1A-1, 1A-2, 1A-3, 1A-4, 1A-5, 1A-6, 1B-1, 5B-2

Cause-specific death rates 2B-2, 2B-3, 2B-4, 2B-5, 2B-6, 2C-6, 2C-7, 2C-8, 2C-9, 2C-12, 2C-13, 2C-16, 2C-17, 2C-20, 2C-21, 2C-24, 2C-25, 5E-11, 5E-13, 5E-20, 5E-26, 5E-28, 5E-30, 5E-32, 5E-34

Crude death rates 2B-6, 5E-13

Fertility rates 1A-1, 1A-2, 1A-3, 1A-4, 1A-5, 1A-6, 1B-1, 5A-1

Natural increase 5B-1

Pregnancy rates 1A-1, 1A-2, 1A-3, 1A-4, 1A-5, 1A-6, 1A-7, 5A-1, 5A-2, 5A-3

**Rectal atresia/stenosis**, (*see also* **Complications of labor/delivery**) 1B-26,

1B-28, 1B-32, 1B-33, 1B-34, 5B-27

**Relapsing fever, Tick** 3A-1, 5F-1, 5F-2

**Renal agenesis**, (*see also* **Complications of labor/delivery**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-27

**Renal disease**, (*see also* **Medical risk factors for pregnancy**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25

**Renal failure**, deaths from 2B-6, 2C-28, 5E-12

**Reportable diseases** (*see* **Diseases, Reportable**)

**Respiratory Distress Syndrome**, deaths from, (*see also* **Certain conditions originating in the perinatal period**) 2C-5, 2C-6, 2C-7, 5E-20

**Respiratory infection, acute upper**, emergency department visits for, 4C-2, 7C-2

**Respiratory system, diseases of**, emergency department visits for, 4C-1, 7C-1

**Respiratory system, diseases of**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**Respiratory therapy**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

**Reyes syndrome** 3A-1, 3A-2, 5F-1, 5F-2

**Rheumatism, excluding back**, emergency department visits for, 4C-2, 7C-2

**Rh sensitization**, (*see also* **Medical risk factors for pregnancy**) 1B-26, 1B-28, 1B-32, 1B-34, 5B-25

**Rocky Mountain Spotted Fever** 3A-1, 3A-2, 5F-1, 5F-2

**Rubella** 3A-1, 3A-2, 6A-1, 5F-1, 5F-2

**Rubella syndrome, congenital** 3A-1, 5F-1, 5F-2

**Rupture of membranes**, (*see also* **Complications of labor/delivery**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-27

**Rural areas** 1B-5, 1B-7, 1B-25, 1B-26, 2B-5, 2C-1, 2C-7, 2C-11, 2C-13, 2C-14, 2C-15, 2C-17, 2C-18, 2C-19, 2C-21, 2C-22, 2C-23, 2C-25, 2C-26, 2D-3, 2D-4, 5B-17

**S**

**Salmonella paratyphi A** 3A-1, 5F-1, 5F-2  
**Salmonella paratyphi B** 3A-1, 5F-1, 5F-2  
**Salmonella paratyphi C** 3A-1, 5F-1, 5F-2  
**Salmonellosis** (except *S. typhi* and *S. paratyphi*)  
 Reported cases 3A-1, 5F-1, 5F-2  
 Deaths from 3A-2  
**Scarlet fever and erysipelas** 2B-6, 2C-28, 5E-12  
**Schizophrenic disorders**, emergency room visits for, 4C-1, 7C-1  
**Schizophrenic disorders**, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1  
**Seizures**, (*see also* **Abnormal conditions of the newborn**) 1B-26, 1B-27, 1B-28, 1B-32, 1B-34, 5B-28  
**Self-harm, intentional**, (*see* **Suicide**)

**Septicemia**, deaths from 2B-1, 2B-2, 2B-4, 2B-5, 2B-6, 2B-11, 2C-11, 2C-27, 2C-28, 2C-29, 5E-11, 5E-12, 5E-13, 5E-14, 5E-25, 5E-33, 5E-34  
**Septicemia**, inpatient discharges with 4A-1, 4A-2, 4A-3, 4A-5, 7A-1  
**Shigellosis**, deaths from 2B-6, 2C-28, 2C-29, 3A-2, 5E-12  
**Shock syndrome, toxic** 5A-1, 3A-2, 5F-1, 5F-2  
**Skin and subcutaneous tissue, diseases of**, emergency room visits for, 4C-1, 7C-1  
**Skin and subcutaneous tissue, diseases of**, inpatient discharges with 4A-1, 4A-2, 4A-3, 4A-5, 7A-1  
**Spina bifida/meningocele**, (*see also* **Congenital anomalies**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 1B-35, 5B-29  
**Spinal disorders**, emergency room visits for, 4C-2, 7C-2  
**Spinal tap**, (inpatient procedure) 4B-1, 4B-2, 4B-3, 4B-4, 7B-1  
**Sprains**, emergency room visits for, 4C-1, 4C-2, 7C-1, 7C-2  
**Stillbirths**, (*see* **Fetal deaths**)  
**Stimulation of labor**, (*see also* **Obstetric procedures**) 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 5B-26  
**Streptococcal-Group A** 3A-1, 5F-1, 5F-2  
**Streptococcal-Group B** 3A-1, 5F-1, 5F-2  
**Streptococcal sore throat, scarlatina and erysipelas**, deaths from 2B-11  
**Streptococcus pneumoniae** 3A-1, 5F-1, 5F-2  
**Stroke**, (*see* **Cerebrovascular disease**)  
**Substance use during pregnancy** 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25, 5B-30  
**Sudden infant death syndrome (SIDS)** 2C-5, 2C-6, 2C-7, 5E-20  
**Suffocation** 6A-4  
**Suicide (intentional self-harm)** as cause of death 2B-1, 2B-2, 2B-3, 2B-4, 2B-5, 2B-6, 2B-11, 2C-8, 2C-9, 2C-11, 2C-12, 2C-13, 2C-15, 2C-16, 2C-17, 2C-19, 2C-20, 2C-21, 2C-23, 2C-28, 2C-29, 2D-3, 2D-4, 6A-4, 5E-11, 5E-12, 5E-13, 5E-14, 5E-25, 5E-26, 5E-27, 5E-28, 5E-29, 5E-30, 5E-31, 5E-32, 5E-33, 5E-34, 6B-4, 9C  
**Suicide (intentional self-harm)**, emergency room visits related to, 4D-1, 7D-1  
**Suicide (intentional self-harm)**, inpatient discharges related to, 4D-2, 7D-2  
**Symptoms, signs, abnormal findings** (as cause of death) 2B-6, 2C-28, 5E-12  
**Symptoms, signs, and ill-defined conditions**, emergency room visits for, 4C-1, 7C-1  
**Syphilis**  
 Congenital 6A-2  
 Early 3B-3, 3B-4, 3B-5  
 Late 3B-5  
 Primary and secondary 3B-1, 6A-2, 6B-2  
 Reported cases 3B-1, 3B-3, 3B-4, 3B-5, 5F-1, 5F-2  
 Deaths from 2B-6, 2C-28, 2C-29, 3B-2, 5E-12

**T**

**Teenage pregnancies**, (*see* **Pregnancies**)  
**Tobacco use during pregnancy**, (*see also* **Births**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25, 5B-30  
**Tocolysis**, (*see also* **Obstetric procedures**) 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-34, 5B-26  
**Transport accidents**, deaths from, 2B-6, 2C-28, 5E-12  
**Trauma complications and unspecified injuries**, emergency room visits for, 4C-1, 7C-1  
**Tuberculosis** 3A-1, 6A-2, 5F-1, 5F-2, 6B-2  
 Deaths from 2B-6, 2C-28, 5E-12  
 Pulmonary 3A-1, 3A-2, 5F-1, 5F-2  
**Tularemia**  
 Deaths from 3A-2  
 Reported cases 3A-1, 5F-1  
**Typhoid fever** 3A-1, 3A-2, 5F-1, 5F-2

**U**

**Ulcer, peptic**, deaths from 2B-6, 2C-28, 2C-29, 5E-12  
**Ultrasound**, (*see also* **Obstetric procedures**) 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-34, 5B-26  
**Ultrasound**, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1  
**Urban areas** 1B-4, 1B-25, 1B-26, 2B-5, 2C-1, 2C-7, 2C-11, 2C-13, 2C-14, 2C-15, 2C-17, 2C-18, 2C-19, 2C-21, 2C-22, 2C-23, 2C-25, 2C-26, 2D-3, 2D-4, 5B-17  
**Uterine bleeding**, (*see also* **Medical risk factors for pregnancy**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25

**V**

**Valley Fever (Coccidioidomycosis)**, (*see also* **Diseases, Reportable**) 3A-1, 3A-2, 5F-1, 5F-2  
**Vancomycin resistant Enterococcus spp.** 3A-1, **Varicella** (*see* **Chickenpox**)  
**Very low-birthweight** 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15, 1B-26, 1B-28, 1B-32  
**Vibrio spp.** 3A-1, 5F-1, 5F-2  
**Viral hepatitis**, deaths from 2B-6, 2C-28, 5E-12  
**Volume depletion**, inpatient discharges with 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

**W**

**White, non-Hispanic residents** 1A-1, 1A-2, 1A-3, 1A-4, 1A-5, 1A-6, 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-20, 1B-22, 1B-23, 1B-24, 1B-25, 1B-26, 1B-28, 1B-30, 1B-31, 1B-32, 1B-33, 1B-34, 1B-35, 1C-4, 1C-5, 1D-1, 1D-2, 1D-4, 1D-5, 2A-1, 2B-3, 2B-4, 2C-4, 2C-11, 2C-15, 2C-19, 2C-23, 2C-27, 2D-1, 2D-2, 2D-3, 2D-4, 3B-5, 3C-3, 5B-7, 5B-8, 5B-10, 5B-15, 5B-18, 5B-22, 5C-5, 5D-5, 5E-6, 5E-18  
**Whooping cough** (*see also* **Pertussis**)  
 Reported cases 3A-1, 5F-1, 5F-2  
 Deaths from 2B-6, 2C-28, 2C-29, 3A-2, 5E-12

**Y**

**Yersiniosis** (except *Y. pestis*) 3A-1, 5F-1, 5F-2