## TABLE 2B-2 (correction) AGE-ADJUSTED\* MORTALITY RATES BY GENDER AND YEAR FOR SELECTED LEADING CAUSES OF DEATH, ARIZONA, 2000-2010

		2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
TOTAL, ALL CAUSES												
	Total	785.6	791.2	796.4	784.0	757.3	772.5	727.3	663.2	659.6	653.2	679.9
	Male	941.1	952.4	964.8	942.8	909.0	925.7	861.6	773.8	777.6	763.9	802.0
	Female	654.6	656.7	658.0	651.5	632.0	644.9	611.8	561.7	554.3	555.1	572.6
CARDIOVASCULAR DISEASE												
	Total	276.1	267.3	265.5	258.4	248.5	245.7	216.4	194.4	194.1	188.8	188.6
	Male	333.1	321.6	321.1	310.3	298.4	298.2	258.3	227.5	235.8	228.1	225.1
	Female	230.0	223.0	221.0	215.8	207.8	203.8	181.0	164.3	158.1	154.9	156.7
DISEASES OF HEART									-			
	Total	206.1	201.7	201.0	197.1	186.7	186.9	166.1	149.5	149.3	145.9	143.3
	Male	259.3	251.3	253.6	246.6	234.2	237.4	207.3	184.0	189.3	183.1	179.8
	Female	163.6	161.5	158.9	157.1	148.5	147.3	131.8	119.4	115.1	114.1	112.2
CEREBROVASCULAR DISEASE				ı			ı					
	Total	51.7	47.7	47.0	43.9	43.6	40.6	34.8	30.6	30.3	28.9	30.7
	Male	50.8	48.3	47.3	42.9	42.7	40.0	33.4	27.7	29.6	28.2	28.9
	Female	51.8	46.8	46.7	43.8	43.6	40.5	35.2	32.2	30.5	29.1	31.3
MALIGNANT NEOPLASMS			[				_ [	ı			_	
	Total	170.4	169.2	168.6	169.4	164.7	161.8	154.7	148.0	145.5	147.2	150.5
	Male 	204.0	199.8	209.1	208.3	201.2	197.1	187.0	176.7	173.8	174.9	180.6
CLIDONI C I OWED DECDIDATORY	Female	145.8	147.5	139.1	141.3	138.6	136.1	131.0	125.6	123.0	125.8	127.1
CHRONIC LOWER RESPIRATORY DISEASE	T-4-1	47.4	47.0	47.0	45.4	44.0	47.0	44.0	20.0	40.4	40.5	40.7
DISEASE	Total	47.4 56.2	46.9	47.0	45.4	41.9	46.8	44.3 50.2	39.2 43.5	42.1	40.5 44.7	42.6
	Male		53.9	57.1	53.8	48.1	51.3	40.1		46.5		48.1
ACCIDENTS (UNINTENTIONAL	Female	41.3	42.2	40.3	40.0	37.7	43.5	40.1	36.0	38.9	37.1	38.6
INJURIES)	Total	41.1	46.6	45.2	44.8	46.0	51.1	51.2	46.6	44.7	43.1	43.8
•	Male	58.0	66.5	62.8	61.6	61.1	68.7	68.5	60.6	59.0	56.6	56.6
	Female	24.9	27.7	28.8	28.8	31.7	34.4	34.3	32.3	30.6	29.8	31.6
MOTOR VEHICLE ACCIDENTS	romaio	21.7	27.7	20.0	20.0	01.7	01.1	01.0	02.0	00.0	27.0	01.0
	Total	17.5	17.2	18.5	18.0	18.1	18.9	19.6	16.2	13.6	11.6	11.1
	Male	24.4	24.2	25.5	24.9	23.9	27.0	26.8	22.2	19.5	16.8	15.9
	Female	10.6	10.1	11.6	11.2	12.5	11.1	12.4	10.1	7.9	6.5	6.4
INFLUENZA AND PNEUMONIA**				9		0		. –			0	
	Total	24.4	21.9	25.0	23.4	20.1	22.3	18.4	12.9	15.6	14.6	10.9
	Male	29.2	26.8	27.8	29.8	25.7	26.6	22.2	14.9	17.7	15.6	12.8
	Female	21.1	18.1	22.9	18.7	16.3	19.2	15.7	11.5	14.1	13.7	9.3

TABLE 2B-2 (continued)

AGE-ADJUSTED\* MORTALITY RATES BY GENDER AND YEAR FOR SELECTED LEADING CAUSES OF DEATH,

ARIZONA, 2000-2010

		2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
ALZHEIMER'S DISEASE**		<u> </u>	<u></u>	<u> </u>	<u> </u>	U.	<u> </u>	<u> </u>	U	<u> </u>	L.	
	Total	21.8	22.5	28.0	32.5	31.3	32.7	33.4	29.6	29.8	29.5	35.1
	Male	17.5	18.4	23.4	26.6	26.4	27.7	27.0	22.8	23.9	24.5	28.5
	Female	24.2	25.1	30.5	35.9	34.2	35.7	47.4	34.1	33.6	32.8	39.3
DIABETES			''	<u>'</u>	'	,	'		'		<u>'</u>	
	Total	19.0	19.9	22.3	20.3	20.7	20.1	18.9	17.0	17.0	15.7	20.1
	Male	21.0	22.9	25.6	23.6	23.4	24.7	22.5	20.5	20.7	18.2	24.9
	Female	17.2	17.5	19.7	17.5	18.4	16.7	15.9	13.9	13.9	13.5	16.0
CHRONIC LIVER DISEASE AND												
	Total	12.8	12.3	12.1	11.5	11.1	12.7	11.4	11.3	11.7	12.2	12.3
	Male	17.7	17.7	15.6	16.2	15.6	17.7	14.3	15.4	15.3	15.4	16.1
	Female	8.3	7.5	8.9	7.1	6.9	8.1	8.5	7.6	8.4	9.1	8.8
NEPHRITIS, NEPHROTIC SYNDROME,												
AND NEPHROSIS**	Total	11.7	12.0	11.3	10.1	11.0	10.4	8.8	7.6	7.1	7.7	7.6
	Male	13.9	15.6	12.9	12.0	13.4	12.7	10.5	8.8	8.6	9.2	9.0
	Female	10.4	9.7	10.4	8.8	9.4	8.8	7.4	6.6	5.9	6.5	6.4
INTENTIONAL SELF-HARM (SUICIDE)												
	Total	14.6	14.9	15.9	14.6	14.9	15.4	15.4	15.4	14.8	16.1	16.7
	Male	24.7	24.6	26.4	24.0	24.1	24.9	24.7	24.4	23.0	24.6	27.1
	Female	5.2	5.6	6.0	5.8	6.3	6.5	6.6	6.7	7.0	8.1	6.7
ASSAULT (HOMICIDE)												
	Total	7.6	8.7	8.6	8.3	8.2	8.2	8.5	7.7	7.1	5.5	6.5
	Male	11.1	13.7	13.9	12.8	13.3	13.5	13.5	11.8	11.2	8.0	10.5
	Female	3.8	3.6	3.0	3.6	2.9	2.6	3.1	3.2	2.7	2.9	2.4
SEPTICEMIA**												
	Total	6.7	7.2	8.3	8.4	8.0	8.4	8.6	8.1	7.9	7.7	7.4
	Male	7.3	8.0	9.6	9.4	9.4	9.7	9.4	8.4	9.4	8.2	10.8
	Female	6.1	6.5	7.2	7.6	7.0	7.5	8.0	7.9	6.7	7.3	5.1

<sup>\*</sup> The number of deaths per 100,000 population adjusted to the 2000 U.S. standard. The cause-of-death titles are according to the Tenth Revision of the International Classification of Diseases (ICD-10). Beginning in 2000, the causes of death are classified by ICD-10. The causes of death for 1998-1999 are classified by the Ninth Revision (ICD-9). The rates for 1998-1999 were revised in this edition of the report for consistency with the newly recomputed mortality rates in Table 2B-3.

<sup>\*\*</sup> The rates for 1998-1999 are comparability-modified. In order to obtain comparability-modified data, the number of deaths or the death rate for a particular cause of death (such as Alzheimer's disease, Nephritis, Influenza and pneumonia, or Septicemia) classified by ICD-9 is adjusted by multiplying it by the comparability ratio for that cause. Comparability ratios are measures of comparison between ICD-9 and ICD-10. Comparability ratio of 1.0 indicates that the same number of deaths would be assigned to a cause-of-death when ICD-9 or ICD-10 was used. The comparability ratios for causes of death other than the four selected above are within 1.0-1.1 range (see *Technical Notes*).