

**TABLE 3B-5
FREQUENCY OF REPORTED CASES, PERCENT DISTRIBUTION AND RATES OF EARLY AND LATE SYPHILIS,
GONORRHEA, CHLAMYDIA AND HERPES BY ETHNICITY, ARIZONA, 2010**

Race/ethnicity	SYPHILIS						GONORRHEA						CHLAMYDIA			HERPES		
	Early			Late			Resistant			Total			Cases	%	Rate	Cases	%	Rate
	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate						
White Non-Hispanic	142.0	35.9	3.8	118.0	23.9	3.2	0.0	0.0	0.0	795.0	24.5	21.5	6459.0	24.0	174.8	684.0	36.9	18.5
Black or African American	36.0	9.1	15.1	58.0	11.8	24.3	0.0	0.0	0.0	731.0	22.5	305.7	2746.0	10.2	1148.5	173.0	9.3	72.4
Hispanic or Latino	141.0	35.6	7.4	211.0	42.8	11.1	0.0	0.0	0.0	986.0	30.3	52.0	9366.0	34.9	494.2	430.0	23.2	22.7
Asian or Pacific Islander	4.0	1.0	2.3	12.0	2.4	7.0	0.0	0.0	0.0	19.0	0.6	11.1	269.0	1.0	157.8	17.0	0.9	10.0
American Indian or Alaska Native	32.0	8.1	12.4	31.0	6.3	12.0	0.0	0.0	0.0	219.0	6.7	85.1	2683.0	10.0	1042.2	196.0	10.6	76.1
Not Specified	41.0	10.4	N/A	63.0	12.8	N/A	0.0	0.0	NA	499.0	15.4	N/A	5338.0	19.9	N/A	353.0	19.1	N/A
Total	396.0	100.0	6.2	493.0	100.0	7.7	0.0	0.0	0.0	3249.0	100.0	50.8	26861.0	100.0	420.2	1853.0	100.0	29.0

Note: Table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2010.

Note: Rates per 100,000 population.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.