

**TABLE 5B-21
DISTRIBUTION OF LOW-BIRTHWEIGHT (LBW) BIRTHS AND LBW RISK BY
NUMBER OF PRENATAL VISITS AND COUNTY OF RESIDENCE, ARIZONA, 2010**

| | | | Prenatal visits | | | | | Total | |
|------------|---------------------------|----------------------|-----------------|------------|------------|-------------|------------|-------|---------|
| | | | No visits | 1-4 visits | 5-8 visits | 9-12 visits | 13+ visits | | Unknown |
| Apache | Births <2,500 grams (LBW) | Count | 4 | 11 | 23 | 26 | 9 | 2 | 75 |
| | | Share of LBW births* | 5.3% | 14.7% | 30.7% | 34.7% | 12.0% | 2.7% | 100.0% |
| | | LBW risk** | 19.0% | 10.6% | 7.4% | 5.4% | 5.1% | 40.0% | 6.8% |
| | Total births | Count | 21 | 104 | 312 | 481 | 176 | 5 | 1,099 |
| Cochise | Births <2,500 grams (LBW) | Count | 10 | 14 | 38 | 40 | 38 | 2 | 142 |
| | | Share of LBW births* | 7.0% | 9.9% | 26.8% | 28.2% | 26.8% | 1.4% | 100.0% |
| | | LBW risk** | 17.9% | 21.9% | 16.9% | 5.5% | 5.4% | 28.6% | 8.0% |
| | Total births | Count | 56 | 64 | 225 | 723 | 706 | 7 | 1,781 |
| Coconino | Births <2,500 grams (LBW) | Count | 1 | 9 | 37 | 54 | 27 | 3 | 131 |
| | | Share of LBW births* | 0.8% | 6.9% | 28.2% | 41.2% | 20.6% | 2.3% | 100.0% |
| | | LBW risk** | 4.5% | 10.5% | 12.8% | 6.3% | 5.6% | 8.1% | 7.4% |
| | Total births | Count | 22 | 86 | 290 | 857 | 483 | 37 | 1,775 |
| Gila | Births <2,500 grams (LBW) | Count | 4 | 8 | 17 | 22 | 8 | 0 | 59 |
| | | Share of LBW births* | 6.8% | 13.6% | 28.8% | 37.3% | 13.6% | 0.0% | 100.0% |
| | | LBW risk** | 33.3% | 19.5% | 9.4% | 7.1% | 6.3% | 0.0% | 8.8% |
| | Total births | Count | 12 | 41 | 181 | 308 | 128 | 0 | 670 |
| Graham | Births <2,500 grams (LBW) | Count | 1 | 4 | 16 | 4 | 2 | 0 | 27 |
| | | Share of LBW births* | 3.7% | 14.8% | 59.3% | 14.8% | 7.4% | 0.0% | 100.0% |
| | | LBW risk** | 20.0% | 6.9% | 8.6% | 1.8% | 3.8% | 0.0% | 5.1% |
| | Total births | Count | 5 | 58 | 186 | 228 | 53 | 0 | 530 |
| Greenlee | Births <2,500 grams (LBW) | Count | 0 | 3 | 3 | 3 | 1 | 0 | 10 |
| | | Share of LBW births* | 0.0% | 30.0% | 30.0% | 30.0% | 10.0% | 0.0% | 100.0% |
| | | LBW risk** | 0.0% | 17.6% | 7.9% | 7.1% | 12.5% | 0.0% | 9.5% |
| | Total births | Count | 0 | 17 | 38 | 42 | 8 | 0 | 105 |
| Maricopa | Births <2,500 grams (LBW) | Count | 103 | 240 | 784 | 1,523 | 1,179 | 22 | 3,851 |
| | | Share of LBW births* | 2.7% | 6.2% | 20.4% | 39.5% | 30.6% | 0.6% | 100.0% |
| | | LBW risk** | 15.0% | 17.5% | 11.8% | 5.7% | 6.4% | 18.6% | 7.1% |
| | Total births | Count | 687 | 1,368 | 6,623 | 26,924 | 18,509 | 118 | 54,236 |
| Mohave | Births <2,500 grams (LBW) | Count | 5 | 5 | 26 | 57 | 47 | 1 | 141 |
| | | Share of LBW births* | 3.5% | 3.5% | 18.4% | 40.4% | 33.3% | 0.7% | 100.0% |
| | | LBW risk** | 19.2% | 10.0% | 9.5% | 6.7% | 5.7% | 33.3% | 7.0% |
| | Total births | Count | 26 | 50 | 275 | 848 | 820 | 3 | 2,022 |
| Navajo | Births <2,500 grams (LBW) | Count | 11 | 15 | 47 | 63 | 18 | 0 | 154 |
| | | Share of LBW births* | 7.1% | 9.7% | 30.5% | 40.9% | 11.7% | 0.0% | 100.0% |
| | | LBW risk** | 31.4% | 9.6% | 9.5% | 8.3% | 6.3% | 0.0% | 8.9% |
| | Total births | Count | 35 | 156 | 497 | 756 | 286 | 7 | 1,737 |
| Pima | Births <2,500 grams (LBW) | Count | 20 | 74 | 266 | 294 | 198 | 1 | 853 |
| | | Share of LBW births* | 2.3% | 8.7% | 31.2% | 34.5% | 23.2% | 0.1% | 100.0% |
| | | LBW risk** | 9.3% | 16.4% | 12.1% | 5.1% | 5.5% | 11.1% | 7.0% |
| | Total births | Count | 215 | 450 | 2,195 | 5,719 | 3,581 | 9 | 12,169 |
| Pinal | Births <2,500 grams (LBW) | Count | 10 | 19 | 79 | 140 | 82 | 0 | 330 |
| | | LBW risk** | 3.0% | 5.8% | 23.9% | 42.4% | 24.8% | 0.0% | 100.0% |
| | | Share of LBW births* | 12.5% | 16.7% | 15.2% | 4.7% | 6.5% | 0.0% | 6.6% |
| | Total births | Count | 80 | 114 | 520 | 3,007 | 1,268 | 1 | 4,990 |
| Santa Cruz | Births <2,500 grams (LBW) | Count | 2 | 8 | 19 | 19 | 10 | 0 | 58 |
| | | Share of LBW births* | 3.4% | 13.8% | 32.8% | 32.8% | 17.2% | 0.0% | 100.0% |
| | | LBW risk** | 10.5% | 9.9% | 8.4% | 6.9% | 11.2% | 0.0% | 8.4% |
| | Total births | Count | 19 | 81 | 227 | 277 | 89 | 0 | 693 |
| Yavapai | Births <2,500 grams (LBW) | Count | 1 | 9 | 27 | 41 | 35 | 2 | 115 |
| | | Share of LBW births* | 0.9% | 7.8% | 23.5% | 35.7% | 30.4% | 1.7% | 100.0% |
| | | LBW risk** | 5.9% | 13.8% | 13.6% | 5.3% | 4.6% | 66.7% | 6.3% |
| | Total births | Count | 17 | 65 | 199 | 771 | 762 | 3 | 1,817 |
| Yuma | Births <2,500 grams (LBW) | Count | 22 | 39 | 70 | 61 | 12 | 0 | 204 |
| | | Share of LBW births* | 10.8% | 19.1% | 34.3% | 29.9% | 5.9% | 0.0% | 100.0% |
| | | LBW risk** | 12.2% | 16.7% | 9.9% | 4.1% | 2.0% | 0.0% | 6.3% |
| | Total births | Count | 181 | 233 | 708 | 1,498 | 609 | 0 | 3,229 |
| La Paz | Births <2,500 grams (LBW) | Count | 1 | 1 | 1 | 2 | 0 | 0 | 5 |
| | | Share of LBW births* | 20.0% | 20.0% | 20.0% | 40.0% | 0.0% | 0.0% | 100.0% |
| | | LBW risk** | 14.3% | 10.0% | 2.1% | 2.8% | 0.0% | 0.0% | 2.5% |
| | Total births | Count | 7 | 10 | 48 | 71 | 62 | 2 | 200 |

* Percent distribution of LBW births by number of prenatal visits.

** Number of LBW births per 100 births in specified prenatal care group.