TABLE 2B-2 (correction) AGE-ADJUSTED* MORTALITY RATES BY GENDER AND YEAR FOR SELECTED LEADING CAUSES OF DEATH, ARIZONA, 2001-2011

		2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
TOTAL, ALL CAUSES				<u> </u>								
	Total	791.2	796.4	784.0	757.3	772.5	727.3	663.2	659.6	653.2	679.9	699.6
	Male	952.4	964.8	942.8	909.0	925.7	861.6	773.8	777.6	763.9	802.0	817.1
	Female	656.7	658.0	651.5	632.0	644.9	611.8	561.7	554.3	555.1	572.6	595.3
CARDIOVASCULAR DISEASE												
	Total	267.3	265.5	258.4	248.5	245.7	216.4	194.4	194.1	188.8	188.6	201.0
	Male	321.6	321.1	310.3	298.4	298.2	258.3	227.5	235.8	228.1	225.1	239.2
	Female	223.0	221.0	215.8	207.8	203.8	181.0	164.3	158.1	154.9	156.7	168.0
DISEASES OF HEART			ı									
	Total	201.7	201.0	197.1	186.7	186.9	166.1	149.5	149.3	145.9	143.3	152.7
	Male	251.3	253.6	246.6	234.2	237.4	207.3	184.0	189.3	183.1	179.8	192.8
	Female	161.5	158.9	157.1	148.5	147.3	131.8	119.4	115.1	114.1	112.2	119.0
CEREBROVASCULAR DISEASE							ı	ı			ı	
	Total	47.7	47.0	43.9	43.6	40.6	34.8	30.6	30.3	28.9	30.7	30.6
	Male	48.3	47.3	42.9	42.7	40.0	33.4	27.7	29.6	28.2	28.9	28.0
	Female	46.8	46.7	43.8	43.6	40.5	35.2	32.2	30.5	29.1	31.3	32.3
MALIGNANT NEOPLASMS	Total	169.2	168.6	169.4	164.7	161.8	154.7	148.0	145.5	147.2	150.5	151.3
	Male	199.2	209.1	208.3	201.2	197.1	187.0	176.7	173.8	174.9	180.5	179.3
	Female	147.5	139.1	141.3	138.6	136.1	131.0	125.6	123.0	125.8	127.1	179.5
CHRONIC LOWER RESPIRATORY	remale	147.5	137.1	141.3	130.0	130.1	131.0	125.0	123.0	125.6	127.1	129.5
DISEASE	Total	46.9	47.0	45.4	41.9	46.8	44.3	39.2	42.1	40.5	42.6	45.9
	Male	53.9	57.1	53.8	48.1	51.3	50.2	43.5	46.5	44.7	48.1	49.7
	Female	42.2	40.3	40.0	37.7	43.5	40.1	36.0	38.9	37.1	38.6	42.8
ACCIDENTS (UNINTENTIONAL												
INJURIES)	Total	46.6	45.2	44.8	46.0	51.1	51.2	46.6	44.7	43.1	43.8	45.5
	Male	66.5	62.8	61.6	61.1	68.7	68.5	60.6	59.0	56.6	56.6	58.3
	Female	27.7	28.8	28.8	31.7	34.4	34.3	32.3	30.6	29.8	31.6	32.6
MOTOR VEHICLE ACCIDENTS												
	Total	17.2	18.5	18.0	18.1	18.9	19.6	16.2	13.6	11.6	11.1	12.1
	Male	24.2	25.5	24.9	23.9	27.0	26.8	22.2	19.5	16.8	15.9	17.6
	Female	10.1	11.6	11.2	12.5	11.1	12.4	10.1	7.9	6.5	6.4	6.9
INFLUENZA AND PNEUMONIA**												
	Total	21.9	25.0	23.4	20.1	22.3	18.4	12.9	15.6	14.6	10.9	9.7
	Male	26.8	27.8	29.8	25.7	26.6	22.2	14.9	17.7	15.6	12.8	11.5
	Female	18.1	22.9	18.7	16.3	19.2	15.7	11.5	14.1	13.7	9.3	8.2

TABLE 2B-2 (continued)

AGE-ADJUSTED* MORTALITY RATES BY GENDER AND YEAR FOR SELECTED LEADING CAUSES OF DEATH,

ARIZONA, 2001-2011

		2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
ALZHEIMER'S DISEASE**		L L	<u></u>	<u> </u>	<u> </u>	U.	<u> </u>	<u> </u>	U	<u> </u>	U	
	Total	22.5	28.0	32.5	31.3	32.7	33.4	29.6	29.8	29.5	35.1	35.2
	Male	18.4	23.4	26.6	26.4	27.7	27.0	22.8	23.9	24.5	28.5	28.9
	Female	25.1	30.5	35.9	34.2	35.7	47.4	34.1	33.6	32.8	39.3	39.0
DIABETES		1	''		'	,			'		<u>'</u>	
	Total	19.9	22.3	20.3	20.7	20.1	18.9	17.0	17.0	15.7	20.1	24.8
	Male	22.9	25.6	23.6	23.4	24.7	22.5	20.5	20.7	18.2	24.9	30.3
	Female	17.5	19.7	17.5	18.4	16.7	15.9	13.9	13.9	13.5	16.0	20.2
CHRONIC LIVER DISEASE AND												
	Total	12.3	12.1	11.5	11.1	12.7	11.4	11.3	11.7	12.2	12.3	13.4
	Male	17.7	15.6	16.2	15.6	17.7	14.3	15.4	15.3	15.4	16.1	18.0
	Female	7.5	8.9	7.1	6.9	8.1	8.5	7.6	8.4	9.1	8.8	9.2
NEPHRITIS, NEPHROTIC SYNDROME,												
AND NEPHROSIS**	Total	12.0	11.3	10.1	11.0	10.4	8.8	7.6	7.1	7.7	7.6	6.0
	Male	15.6	12.9	12.0	13.4	12.7	10.5	8.8	8.6	9.2	9.0	7.2
	Female	9.7	10.4	8.8	9.4	8.8	7.4	6.6	5.9	6.5	6.4	5.1
INTENTIONAL SELF-HARM (SUICIDE)												
	Total	14.9	15.9	14.6	14.9	15.4	15.4	15.4	14.8	16.1	16.7	17.2
	Male	24.6	26.4	24.0	24.1	24.9	24.7	24.4	23.0	24.6	27.1	27.4
	Female	5.6	6.0	5.8	6.3	6.5	6.6	6.7	7.0	8.1	6.7	7.5
ASSAULT (HOMICIDE)												
	Total	8.7	8.6	8.3	8.2	8.2	8.5	7.7	7.1	5.5	6.5	6.1
	Male	13.7	13.9	12.8	13.3	13.5	13.5	11.8	11.2	8.0	10.5	9.6
	Female	3.6	3.0	3.6	2.9	2.6	3.1	3.2	2.7	2.9	2.4	2.6
SEPTICEMIA**												
	Total	7.2	8.3	8.4	8.0	8.4	8.6	8.1	7.9	7.7	7.4	4.7
	Male	8.0	9.6	9.4	9.4	9.7	9.4	8.4	9.4	8.2	10.8	5.4
	Female	6.5	7.2	7.6	7.0	7.5	8.0	7.9	6.7	7.3	5.1	4.0

^{*} The number of deaths per 100,000 population adjusted to the 2000 U.S. standard. The cause-of-death titles are according to the Tenth Revision of the International Classification of Diseases (ICD-10). Beginning in 2000, the causes of death are classified by ICD-10. The causes of death for 1998-1999 are classified by the Ninth Revision (ICD-9). The rates for 1998-1999 were revised in this edition of the report for consistency with the newly recomputed mortality rates in Table 2B-3.

^{**} The rates for 1998-1999 are comparability-modified. In order to obtain comparability-modified data, the number of deaths or the death rate for a particular cause of death (such as Alzheimer's disease, Nephritis, Influenza and pneumonia, or Septicemia) classified by ICD-9 is adjusted by multiplying it by the comparability ratio for that cause. Comparability ratios are measures of comparison between ICD-9 and ICD-10. Comparability ratio of 1.0 indicates that the same number of deaths would be assigned to a cause-of-death when ICD-9 or ICD-10 was used. The comparability ratios for causes of death other than the four selected above are within 1.0-1.1 range (see *Technical Notes*).