

**TABLE 3B-5
FREQUENCY OF REPORTED CASES, PERCENT DISTRIBUTION AND RATES OF EARLY AND LATE SYPHILIS,
GONORRHEA, CHLAMYDIA, AND HERPES BY ETHNICITY, ARIZONA, 2011**

Race/ethnicity	SYPHILIS						GONORRHEA						CHLAMYDIA			HERPES		
	Early			Late			Resistant			Total			Cases	%	Rate	Cases	%	Rate
	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate						
White Non-Hispanic	176	38.2	4.8	96	22.3	2.6	0	0.0	0.0	1,067	23.4	28.9	6,905	23.6	186.8	832	38.6	22.5
Black or African American	57	12.4	23.8	60	13.9	25.1	0	0.0	0.0	994	21.8	415.7	2,741	9.4	1146.4	200	9.3	83.6
Hispanic or Latino	195	42.3	10.3	220	51.0	11.6	0	0.0	0.0	1,271	27.8	67.1	10,102	34.5	533.0	569	26.4	30.0
Asian or Pacific Islander	3	0.7	1.8	4	0.9	2.3	0	0.0	0.0	34	0.7	19.9	226	0.8	132.5	14	0.6	8.2
American Indian or Alaska Native	27	5.9	10.5	37	8.6	14.4	0	0.0	0.0	419	9.2	162.8	2,908	9.9	1129.6	176	8.2	68.4
Not Specified	3	0.7	N/A	14	3.2	N/A	0	0.0	NA	779	17.1	N/A	6,369	21.8	N/A	364	16.9	N/A
Total	461	100.0	7.2	431	100.0	6.7	0	0.0	0.0	4,564	100.0	71.4	29,251	100.0	457.6	2,155	100.0	33.7

Note: Table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2010.

Note: Rates per 100,000 population.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.