

**TABLE 3B-3
FREQUENCY OF REPORTED CASES OF GONORRHEA, CHLAMYDIA, AND EARLY SYPHILIS
BY AGE AND GENDER, ARIZONA, 2015**

Age group	GONORRHEA				CHLAMYDIA				EARLY SYPHILIS			
	Males	Females	Unknown or Transgender	Total	Males	Females	Unknown or Transgender	Total	Males	Females	Unknown or Transgender	Total
0-4	0	0	0	0	0	0	0	0	0	0	0	0
5-9	0	*	0	0†	0	0	0	0	0	0	0	0
10-14	6	30	0	36	18	131	0	149	0	0	0	0
15-19	510	751	*	1,264	1,687	5,952	6	7,645	36	8	0	44
20-24	1,365	1,150	*	2,517	3,637	8,958	17	12,612	151	30	0	181
25-29	1,040	761	*	1,805	2,190	3,949	13	6,152	156	20	0	176
30-34	696	395	0	1,091	1,124	1,754	*	2,882	127	12	0	139
35-39	409	208	*	618	572	854	*	1,427	109	14	0	123
40-44	275	100	*	376	344	410	*	755	73	*	0	80†
45-49	183	68	0	251	242	205	*	448	84	*	0	90†
50-54	130	35	0	165	128	111	0	239	69	*	0	70†
55-59	66	15	0	81	62	52	0	114	33	0	0	33
60-64	35	*	0	40†	36	18	0	54	14	0	0	14
65-over	23	*	0	30†	28	*	*	30†	8	0	0	8
Total	4,738	3,520†	10†	8,270†	10,068	22,400†	40†	32,511	860	90†	0	952

Notes: * Cell suppressed due to non-zero count less than 6; † Sum rounded to nearest tens unit due to non-zero addend less than 6; since 2005, the table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.