

**TABLE 3B-3
FREQUENCY OF REPORTED CASES OF GONORRHEA, CHLAMYDIA, AND EARLY SYPHILIS
BY AGE AND GENDER, ARIZONA, 2016**

Age group	GONORRHEA				CHLAMYDIA				EARLY SYPHILIS			
	Males	Females	Unknown or Transgender	Total	Males	Females	Unknown or Transgender	Total	Males	Females	Unknown or Transgender	Total
0-4	*	*	0	10†	0	6	0	6	0	0	0	0
5-9	0	*	0	0†	0	*	0	0†	0	0	0	0
10-14	10	38	0	48	26	166	*	193	0	0	0	0
15-19	659	856	0	1,515	1,967	6,385	*	8,356	35	7	0	42
20-24	1,633	1,388	*	3,022	3,898	9,145	*	13,046	111	21	0	132
25-29	1,333	894	*	2,228	2,352	4,330	*	6,683	140	7	0	147
30-34	837	540	*	1,378	1,277	1,838	*	3,117	97	15	0	112
35-39	514	248	*	763	715	894	*	1,610	58	9	0	67
40-44	371	170	0	541	388	479	0	867	42	10	0	52
45-49	282	94	0	376	280	214	*	495	59	*	0	60†
50-54	209	51	0	260	165	145	0	310	56	6	0	62
55-59	93	19	0	112	77	59	0	136	20	0	0	20
60-64	41	8	0	49	41	15	0	56	17	0	0	17
65-over	28	*	0	30†	29	16	0	45	6	*	0	10†
Total	6,010†	4,320†	0†	10,330†	11,215	23,700†	10†	34,920†	641	80†	0	721

Notes: * Cell suppressed due to non-zero count less than 6; † Sum rounded to nearest tens unit due to non-zero addend less than 6; since 2005, the table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.