

**TABLE 3B-4
RATES^a OF REPORTED CASES OF GONORRHEA, CHLAMYDIA, AND EARLY SYPHILIS
BY AGE AND GENDER, ARIZONA, 2016**

Age group	GONORRHEA			CHLAMYDIA			EARLY SYPHILIS		
	Males	Females	Total	Males	Females	Total	Males	Females	Total
0-4	**	**	**	0.0	2.9	1.4	0.0	0.0	0.0
5-9	0.0	**	**	0.0	**	**	0.0	0.0	0.0
10-14	4.3	16.9	10.5	11.2	73.9	42.2	0.0	0.0	0.0
15-19	279.0	380.8	328.6	832.8	2840.4	1812.7	19.1	7.6	13.4
20-24	646.3	591.7	620.2	1542.7	3898.5	2677.4	64.1	17.9	41.9
25-29	550.4	398.5	477.6	971.2	1930.1	1432.5	84.6	9.8	48.7
30-34	365.1	247.9	308.2	557.0	843.6	697.1	65.0	11.0	38.7
35-39	240.7	118.3	180.3	334.9	426.3	380.4	52.9	10.5	31.9
40-44	176.8	81.5	129.3	184.9	229.6	207.2	44.8	6.7	25.8
45-49	136.6	45.6	91.1	135.7	103.8	120.0	54.7	**	29.1
50-54	98.5	23.3	60.3	77.8	66.2	71.9	42.9	5.5	23.9
55-59	45.7	8.6	26.4	37.8	26.7	32.0	18.2	0.0	8.7
60-64	22.4	3.9	12.6	22.4	7.3	14.4	15.3	0.0	7.2
65-over	5.4	**	2.9	5.6	2.6	4.0	2.1	**	1.1
Total	177.2	125.6	151.3	330.7	689.5	511.5	30.9	4.7	17.7

Notes: ** Cell suppressed due to rate/ratio/percent based on non-zero count less than 6; ^a Number of cases per 100,000 population; table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2016; denominators for unknown or transgender category are not available; rates per 100,000 population.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.