## **Risk Profile of Asians or Pacific Islanders**

(percent above or below the State mean)

	BETTER THAN AVERAGE	WORSE THAN AVERAGE
9.Mortality of infants from low birthweight	-100.0	
6.Drug-induced deaths	-84.8	
2.Incidence of gonorrhea	-82.2	
15.HIV disease 11.Prostate cancer	-79.2	
4.Incidence of genital herpes		
2.Pregnancies among females 14 and younger		
3.Incidence of chlamydia	-71.7	
2.Mortality of adolescents 15-19 years	-69.3	
5.Chronic lower respiratory diseases	-61.8	
14.Chronic liver disease and cirrhosis 1.Unmarried mothers		
2.Tobacco use during pregnancy		
3.Pregnancies among females 15-17 years		
3.No prenatal care		
<ol><li>Mortality of young adults 20-44 years</li></ol>	-56.5	
7.Lung cancer	-54.2	
4.Pregnancies among females 18-19 years	-54.1	
4.Mortality of middle-aged adults 45-64 years	<u>-53.5</u>	
5.Injury by firearms 1.Pregnancies among females 19 and younger		
3.Alcohol use during pregnancy	-50.5	·
1.Incidence of early syphilis		
2.Fewer than five prenatal visits	-46.2	
5.Incidence of AIDS	-46.1	
2.Motor vehicle-related injuries	-46.0	
1.No care in the 1st trimester	-38.6	
9.Assault (homicide) 7.Alcohol-induced deaths		
1.Incidence of hyaline membrane disease		
1.Unintentional injuries		
6.Malignant neoplasm (cancer)		
10.Neonatal mortality	-27.	
16.Septicemia	-26.	
8.Intentional self-harm (suicide)	-24.	0
10.Weight gain of less than 21 pounds	-23	
3.Coronary heart disease		.1
8.Pregnancy-associated hypertension 7.Infant mortality		
12.Influenza and pneumonia		.1
6.Anemia during pregnancy		.0
10.Colorectal cancer		.8
20.Total mortality from all causes		.5 •
1.Mortality of children 1-14 years	-17	
8.Breast cancer		5.4 🔳
2.Diseases of the heart		4.5 📕
11.Postneonatal mortality		2.5
18.Nephritis, nephrotic syndrome and nephrosis 13.Diabetes		1.5
17.Alzheimer's disease		1.4 ■ -7.6 □
2.Incidence of newborn intensive care		-5.3
3.Born too early (preterm or < 37 weeks)		-4.7
1.Cardiovascular disease		-1.1
9.Genital herpes during pregnancy		0.0
5.Born too small: <1,500 grams	L	0.0
19.Essential (primary) hypertension		
2.Median age at death among males 1.Median age at death from all causes		
3.Median age at death among females		<mark>6.5</mark>
4.Born too small: <2,500 grams		• • • • • • • • • • • • • • • • • • •
Incidence of congenital anomalies		
.Deaths before expected years of life reached		23.5
Chronic hypertension		<b>2</b> 5.0
.Dysfunctional labor		25.0
.Fall-related injuries	L	25.0
. Diabetes during pregnancy . Cerebrovascular disease	·	
4.Cerebrovascular disease 9.Cervical cancer		59.1
4.Drowning mortality		
3.Sudden Infant Death Syndrome (SIDS)		128.6
	I I 5 0 0 .0	I I 0.0
-	0.000	0.0
	- 2 5 0 .0	AVER AGE 2 5 0 .0