**Risk Profile of Hispanic or Latino** (percent above or below the State mean)

	BETTER THAN AVERAGE		WORSE THAN AVERAGE	
2-2. Tobacco use during pregnancy		-62.5		
2-3. Alcohol use during pregnancy		-60.0		
5-5. Chronic lower respiratory diseases		-59.5		
6-8. Intentional self-harm (suicide)		-44.1		
6-6. Drug-induced deaths		-41.1		
2-9. Genital herpes during pregnancy 5-7. Lung cancer		-40.0		
9-4. Incidence of genital herpes		-32.5		
2-6. Anemia during pregnancy		-27.8 📕		
2-4. Chronic hypertension		-25.0 📕		
2-5. Dysfunctional labor		-25.0		
2-8. Pregnancy-associated hypertension		-25.0		
7-3. Mortality of young adults 20-44 years		-23.6		
4-1. Incidence of hyaline membrane disease 6-5. Injury by firearms		-18.8		
6-1. Unintentional injuries		-18.1		
7-4. Mortality of middle-aged adults 45-64 years		-17.2 📕		
6-7. Alcohol-induced deaths		-15.8		
4-11. Postneon at al mortality				
5-8. Breast cancer				
5-6. Malignant neoplasm (cancer) 4-8. Sudden Infant Death Syndrome (SIDS)	F	<u>-12.8</u> _ -11.4		
5-2. Diseases of the heart	+	-10.6		
5-3. Coronary heart disease		-10.5		
5-17. Alzheimer's disease		-8.8		
6-2. Motor vehicle-related injuries		-8.6		
4-5. Born too small: <1,500 grams		-8.3		
4-4. Born too small: < 2,500 grams				
6-4. Drowning mortality 4-2. Incidence of newborn intensive care		-6.5		
9-2. Incidence of gonorrhea		-5.9		
5-20. Total mortality from all causes		-5.9		
4-7. Infant mortality		5.1		
6-3. Fall-related injuries				
5-1. Cardiovascular disease		-4.8		
4-3. Born too early (preterm or < 37 weeks) 4-10. Neonatal mortality		-3.0	0.0	
4-6. Incidence of congenital anomalies			0.0	
2-7. Diabetes during pregnancy			2.6	
5-16. Septicemia			<u>3.9</u>	
5-12. Influenza and pneumonia			6.2	
5-10. Colorectal cancer			7.0 8.4	
7-1. Mortality of children 1-14 years 5-11. Prostate cancer			8.7	
8-3. Median age at death among females			10.0	
7-2. Mortality of adolescents 15-19 years			11.3	
8-1. Median age at death from all causes			11.8	
8-2. Median age at death among males			13.7	
5-9. Cervical cancer			<u>16.7</u> 17.0	
5-4. Cerebrovascular disease 4-9. Mortality of infants from low birthweight			18.5	
9-3. Incidence of chlamydia			19.6	
2-10. Weight gain of less than 21 pounds	L		23.5	
2-1. Unmarried mothers	L		25.9	
9-5. Incidence of HIV/AIDS	F+		<u> </u>	
8-4. Deaths before expected years of life reached			31.1	
3-1. No care in the 1st trimester 5-15. HIV disease	F		<b>3</b> 1.3	
5-19. Essential (primary) hypertension			32.4	
3-2. Fewer than five prenatal visits	L		39.6	
9-1. Incidence of early syphilis			43.1	
6-9. Assault (homicide)	F+		<u>47.3</u> 53.3	
5-14. Chronic liver disease and cirrhosis 1-1. Pregnancies among females 19 and younger	+		55.4	
1-1. Pregnancies among females 19 and younger 1-2. Pregnancies among females 14 and younger			57.1	
1-4. Pregnancies among females 18-19 years	L		60.2	
3-3. No prenatal care	L		61.1	
5-13. Diabetes			79.0	
1-3. Pregnancies among females 15-17 years	F			
5-18. Nephritis, nephrotic syndrome and nephrosis	+		00.1	
		2	0	
	0.0 -250.0 AVERAGE 250.0 FOR ALL RESIDENTS			0.0