

RISK PROFILE OF AMERICAN INDIANS IN ARIZONA:
AN OVERVIEW OF SELECTED HEALTH PROBLEMS AND HEALTH RISKS
AMONG AMERICAN INDIANS AND ALL RESIDENTS OF THE STATE IN 2021

The risk profile presented on the following page summarizes how American Indians as a group compared in 2021 to the state average on 66 measures of health status. The 66 indicators used in the profile are individual measures of the various dimensions or facets of health status. The profile uses the percentages a racial/ethnic group is above or below the statewide average on each of the specific measures. In order to represent the level of each indicator for American Indians in comparison to the overall state level, the following formula was used:*

The above formula produces a score of 0.0 for a group with the same value as the statewide average. A negative value indicates better than average standing on an indicator. A positive value indicates worse than average standing on an indicator*. The profile visually describes the better-than-average and worse-than-average clusters of health status characteristics for American Indian residents of Arizona.

Following the graphic profile is a detailed description of the 66 indicators. These indicators are organized around 9 major subject areas:

(1) Teen Pregnancies, (2) Maternal Lifestyle and Health Characteristics, (3) Utilization of Prenatal Care Services, (4) Newborn's Health, (5) Mortality from Natural Causes (chronic or infectious diseases), (6) Mortality from Injuries, (7) Age-Specific Mortality, (8) Premature Mortality and (9) Reportable Diseases.

IN 2021, AMERICAN INDIAN RESIDENTS OF ARIZONA:

- ranked better than average on a number of health indicators including: mortality rates for several chronic diseases including breast cancer, lung cancer, chronic lower respiratory disease, Alzheimer's disease, malignant neoplasms, and prostate cancer;
- ranked worse than the statewide average on 53 of 66 health indicators;
- had high mortality from alcohol-induced causes, chronic liver disease and cirrhosis, COVID-19, septicemia, diabetes, assault,
 influenza and pneumonia, motor vehicle accidents, early syphilis, and unintentional injuries;
- ranked worse on measures of maternal lifestyle and health, as well as in utilization of prenatal care;
- on average were 15 years younger at time of death compared to all racial/ethnic groups.

For consistency, the signs + and - for scores indicating the median age at death as higher or lower than the statewide average were reversed (indicators 8-1, 8-2, 8-3). Otherwise, a lower-than-average median age at death would signify a better than average standing on these indicators. A lower-than-average median age at death obviously reflects a worse off life expectancy.

^{*} The comparative risk profiles for American Indian or Alaska Native, White non-Hispanic, Hispanic or Latino, Black or African American, and Asian or Pacific Islander residents of Arizona were developed for a report "Differences in the Health Status among Racial/Ethnic Groups". This publication is available online at http://pub.azdhs.gov/health-stats/report/dhsag/index.php.

2021 RISK PROFILE OF AMERICAN INDIANS

(Percent above or below the average rate or ratio of all Arizona residents in 2021

AVERAGE FOR ALL ARIZONANS

BETTER THAN AVERAGE	WORSE THAN AVERAGE	
-72.7		4-8.Mortality of infants from low birthweight/short ge
-69.4		5-7.Lung cancer
-56.		5-5.Chronic lower respiratory diseases 5-10.Colorectal cancer
-33		5-10.Colorectal cancer 5-17.Alzheimer's disease
-27		
-24		5-11.Prostate cancer 4-9.Neonatal mortality
-10		
	7	5-3.Coronary heart disease
	.2	5-8.Breast cancer (female)
	.5	5-2.Diseases of the heart
	.2	5-6.Malignant neoplasm (cancer)
	.3	4-3.Born too small: < 2,500 grams
	.0	4-4.Born too small: <1,500 grams
	4	2-6.Gestational weight gain - Inadequate
	.8	4-6.Infant mortality
	.6	5-1.Cardiovascular disease
	1	2-7.Gestational weight gain - Excessive
9	.4	2-2.Tobacco use during pregnancy
1	3.9	4-1.Incidence of newborn intensive care
	8.2	8-3.Median age at death among females
	0.0	4-5.Incidence of congenital anomalies (birth defects)
	0.1	5-4.Cerebrovascular disease
2	0.3	8-1.Median age at death from all causes
2	2.2	8-2.Median age at death among males
	4.0	4-2.Born too early (preterm or < 37 weeks)
	30.6	9-4.Incidence of HIV/AIDS
	38.9	9-3.Incidence of chlamydia
	42.7	1-4.Pregnancies among females 18-19 years
	45.4	6-5.Injury by firearms
	49.5	3-1.No care in the 1st trimester
	50.0	4-7.Sudden Infant Death Syndrome (SIDS)
	50.1	8-4.Deaths before expected years of life reached
	52.9	4-10.Post neonatal mortality
	53.8	1-1.Pregnancies among females 19 and younger
	55.8	6-3.Fall-related injuries
	58.9	5-19.Essential (primary) hypertension
	62.5	6-4.Drowning mortality
	62.6	2-5-Gestational hypertension
		9-2.Incidence of gonorrhea
	62.9 73.7	2-3.Pre-existing hypertension
		2-3.Pre-existing ny pertension 2-1.Unmarried mothers
	75.9	
	79.1	6-6.Drug-induced deaths
	93.1	5-21.Total mortality from all causes
	96.0	2-4.Gestational diabetes
	100.0	5-9.Cervical cancer
	104.6	6-8.Intentional self-harm (suicide)
	116.0	3-3.No prenatal care
	116.4	7-2.Mortality of adolescents 15-19 years
	118.1	1-3.Pregnancies among females 15-17 years
	133.3	1-2.Pregnancies among females 14 and younger
	133.3	3-2.Fewer than five prenatal visits
	149.2	7-4.Mortality of middle-aged adults 45-64 years
	153.5	7-1.Mortality of children 1-14 years
	153.8	5-15.HIV disease
	154.9	5-18.Nephritis, nephrotic syndrome and nephrosis
	165.9	6-1.Unintentional injuries
	169.7	5-12.Influenza and pneumonia
	179.6	5-16.Septicemia
	188.1	5-20.COVID-19
	192.6	6-9.Assault (homicide)
	256.0	6-2.Motor vehicle-related injuries
		9-1.Incidence of early syphilis
	262.0	9-1.Incidence or early syphilis 5-13.Diabetes
	264.2	
	285.6	7-3.Mortality of young adults 20-44 years 5-14.Chronic liver disease and cirrhosis
	574.5	J-14-Cifonic liver disease and cirrhosis
	644.9	6-7.Alcohol-induced deaths

1. TEEN PREGNANCIES

(Number of pregnancies per 1,000 females in specified age group)

Figure Number	Indicator	2021 rate for American Indians	Percent above or below the average rate for all groups	Average rate for all groups in 2021	Definitions
1-1	Pregnancies among females 19 and younger	16.3	53.8	10.6	The sum of live births, fetal deaths, and abortions per 1,000 females aged 19 or younger.
1-2	Pregnancies among females 14 and younger	0.7	133.3	0.3	The sum of live births, fetal deaths, and abortions per 1,000 females aged 14 or younger.
1-3	Pregnancies among females 15-17 years old	18.1	118.1	8.3	The sum of live births, fetal deaths, and abortions per 1,000 females aged 15-17 years.
1-4	Pregnancies among females 18-19 years old	56.1	42.7	39.3	The sum of live births, fetal deaths, and abortions per 1,000 females aged 18-19 years.

2. MATERNAL LIFESTYLE AND HEALTH CHARACTERISTICS

(All rates are per 100 live births)

Figure Number	Indicator	2021 rate for American Indians	Percent above or below the average rate for all groups	Average rate for all groups in 2021	Definitions
2-1	Unmarried mothers	79.5	75.9	45.2	The number of births to unmarried mothers per 100 births.
2-2	Tobacco use during pregnancy	3.5	9.4	3.2	The number of mothers giving birth who self-reported tobacco use during pregnancy per 100 births.
2-3	Pre-existing hypertension	3.3	73.7	1.9	Blood pressure persistently greater than 140/90, diagnosed prior to onset of pregnancy or before the 20th week of gestation (the number of cases per 100 births).
2-4	Gestational diabetes	19.4	96.0	9.9	Metabolic disorder characterized by excessive discharge of urine and persistent thirst; includes juvenile onset, adult onset, and gestational diabetes during pregnancy (the number of cases per 100 births).
2-5	Gestational hypertension	14.8	62.6	9.1	An increase in blood pressure of at least 30 mm hg systolic or 15 mm hg diastolic on two measurements taken 6 hours apart after the 20th week of gestation (the number of cases per 100 births).
2-6	Gestational weight gain - Inadequate	22.4	0.4	22.3	The number of women giving birth whose gestational weight gain was inadequate in relation to pre-pregnancy body mass index (BMI).
2-7	Gestational weight gain - Excessive	49.4	5.1	47.0	The number of women giving birth whose gestational weight gain was excessive in relation to pre-pregnancy body mass index (BMI).

3. UTILIZATION OF PRENATAL CARE SERVICES

(Per 100 live births)

Figure Number	Indicator	2021 rate for American Indians	Percent above or below the average rate for all groups	Average rate for all groups in 2021	Definitions
3-1	No care in the 1 st trimester	42.3	49.5	28.3	The number of mothers giving birth who received no prenatal care in the $1^{\rm st}$ trimester of pregnancy per 100 births.
3-2	Fewer than five prenatal visits	16.8	133.3		The number of mothers giving birth who had fewer than five prenatal visits during pregnancy per 100 births.
3-3	No prenatal care	5.4	116.0	2.5	The number of mothers giving birth who received no prenatal care during pregnancy per 100 births.

4. NEWBORN'S HEALTH

(Infant, neonatal, and postneonatal rates are per 1,000 live births. All other indicators are expressed per 100 live births)

Figure Number	Indicator	2021 rate for American Indians	Percent above or below the average rate for all groups	Average rate for all groups in 2021	Definitions
4-1	Incidence of newborn intensive care	9.0	13.9	7.9	The number of babies admitted to Newborn Intensive Care units per 100 births.
4-2	Born too early (preterm or < 37 weeks)	12.4	24.0	10.0	The number of births before 37 weeks of gestation per 100 births. Preterm is any neonate whose birth occurs through the end of the last day of the 37 th week (259th day), following onset of the last menstrual period.
4-3	Born too small: low-birthweight (<2,500 grams)	7.8	-1.3	7.9	The number of newborns weighing less than 2,500 grams per 100 births. Low-birthweight is any neonate weighing less than 2,500 grams at birth (less than 5 pounds 8 ounces).
4-4	Born too small: very low-birthweight (<1,500 grams)	1.3	0.0	1.3	The number of newborns weighing less than 1,500 grams per 100 births. Very low-birthweight is any neonate whose weight at birth is 1,500 grams or less (less than 3 pounds 5 ounces).
4-5	Incidence of congenital anomalies (birth defects)	0.6	20.0	0.5	The numbers of babies born with birth defects per 100 births.
4-6	Infant mortality	5.6	1.8	5.5	The number of infant deaths per 1,000 live births. Infant death is any death at any time from birth up to, but not including, the first year of age (364 days, 23 hours, 59 minutes from the moment of birth).
4-7	Sudden Infant Death Syndrome (SIDS)	0.3	50.0	0.2	The number of infant deaths from Sudden Infant Death Syndrome (ICD-10 code R95) per 1,000 live births.
4-8	Mortality of infants from low birthweight/short gestation	0.3	-72.7	1.1	The number of infant deaths from low-birthweight/short gestation (ICD-10 code P07) per 1,000 live births.
4-9	Neonatal mortality	3.1	-18.4	3.8	The number of neonatal deaths per 1,000 live births. Neonatal death is death of a liveborn neonate before the neonate becomes 28 days old (up to and including 27 days, 23 hours, 59 minutes from the moment of birth).
4-10	Postneonatal mortality	2.6	52.9	1.7	The number of postneonatal deaths per 1,000 live births. Postneonatal death is any death of a liveborn infant at least 28 days of age but less than one year of age.

5. MORTALITY FROM NATURAL CAUSES (chronic or infectious diseases) (Number of deaths per 100,000 population age-adjusted to 2000 standard. The causes of death are classified by the Tenth Revision of the International Classification of Diseases or ICD-10)

Figure Number	Indicator	2021 rate for American Indians	Percent above or below the average rate for all groups	Average rate for all groups in 2021	Definitions
5-1	Cardiovascular disease	210.8	3.6	203.4	The number of deaths from cardiovascular diseases (ICD-10 codes I00-I78) per 100,000 age-adjusted population.
5-2	Diseases of the heart	145.9	-3.5	151.2	The number of deaths from disease of the heart (ICD-10 codes I00-I09, I11, I13, I20-I51) per 100,000 age-adjusted population.
5-3	Coronary heart disease	97.6	-9.7	108.1	The number of deaths from coronary heart disease (ICD-10 codes I200-I259) per 100,000 age-adjusted population.
5-4	Cerebrovascular disease	41.3	20.1	34.4	The number of deaths from cerebrovascular disease (ICD-10 codes I60-I69) per 100,000 age-adjusted population.
5-5	Chronic lower respiratory diseases	15.2	-56.9	35.3	The number of deaths from chronic lower respiratory diseases (ICD-10 codes J40–J47) per 100,000 age-adjusted population.
5-6	Malignant neoplasm (cancer)	128.9	-2.2	131.8	The number of deaths from cancer (ICD-10 codes C00-C97) per 100,000 age-adjusted population.
5-7	Lung cancer	7.9	-69.4	25.8	The number of deaths from lung cancer (ICD-10 codes C33-C34) per 100,000 age- adjusted population.
5-8	Breast cancer	18.3	-6.2	19.5	The number of deaths from breast cancer (ICD-10 code C50) per 100,000 age-adjusted female population.
5-9	Cervical cancer	5.0	100.0	2.5	The number of deaths from cervical cancer (ICD-10 code C53) per 100,000 age- adjusted female population.
5-10	Colorectal cancer	8.2	-33.9	12.4	The number of deaths from colorectal cancer (ICD-10 codes $C18-C21$) per $100,000$ age-adjusted population.
5-11	Prostate cancer	13.0	-24.4	17.2	The number of deaths from prostate cancer (ICD-10 code C61) per 100,000 age-adjusted male population.
5-12	Influenza and pneumonia	26.7	169.7	9.9	The number of deaths from influenza and pneumonia (ICD-10 codes J10-J18) per 100,000 age-adjusted population.
5-13	Diabetes	97.6	264.2	26.8	The number of deaths from diabetes (ICD-10 codes E10-E14) per 100,000 age-adjusted population.
5-14	Chronic liver disease and cirrhosis	145.7	574.5	21.6	The number of deaths from chronic liver disease and cirrhosis (ICD-10 codes K70, K73–K74) per 100,000 age-adjusted population.
5-15	HIV disease	3.3	153.8	1.3	The number of deaths from HIV disease (ICD-10 codes B20-B24) per 100,000 age- adjusted population.
5-16	Septicemia	15.1	179.6	5.4	The number of deaths from septicemia (ICD-10 codes A40-A41) per 100,000 age- adjusted population.
5-17	Alzheimer's disease	20.6	-27.2	28.3	The number of deaths from Alzheimer's disease (ICD-10 code G30) per 100,000 age- adjusted population.
5-18	Nephritis, nephrotic syndrome and nephrosis (kidney disease)	23.2	154.9	9.1	The number of deaths from nephritis, nephritic syndrome, and nephrosis (ICD-10 codes N00–N07, N17–N19,N25–N27) per 100,000 age-adjusted population.
5-19	Essential (primary) hypertension	19.7	58.9	12.4	The number of deaths from essential (primary) hypertension (ICD-10 codes I10, I12) per 100,000 age-adjusted population.
5-20	COVID-19	394.4	188.1	136.9	The number of deaths from COVID-19 (ICD-10 code U071) per 100,000 age-adjusted population.
5-21	Total mortality from all causes	1704.7	93.1	882.6	Total number of deaths from all causes per 100,000 age-adjusted population.

6. MORTALITY FROM INJURIES

(Number of deaths per 100,000 population age-adjusted to 2000 standard. The causes of death are classified by the Tenth Revision of the International Classification of Diseases or ICD-10. Drug-induced deaths include accidental, suicidal, and homicidal poisoning by drugs, but also disorders due to use of psychoactive substances. Alcohol-induced deaths include accidental, suicidal, and homicidal by drugs, but also disorders due to use of alcohol)

Figure Number	Indicator	2021 rate for American Indians	Percent above or below the average rate for all groups	Average rate for all groups in 2021	Definitions
6-1	Unintentional injuries	206.9	165.9	77.8	The number of deaths from unintentional injuries in accidents (ICD-10 codes V01–X59, Y85–Y86) per 100,000 age-adjusted population.
6-2	Motor vehicle-related injuries	65.5	256.0	18.4	V80.3-V80.5,V81.0-V81.1,V82.0-V82.1, V83-V86, V87.0-V87.8, V88.0-V88.8, V89.0, V89.2) per 100.000 age-adjusted population.
6-3	Fall-related injuries	18.7	55.8	12.0	The number of deaths from fall-related injuries (ICD-10 codes W00-W19) per 100,000 age-adjusted population.
6-4	Drowning mortality	2.6	62.5	1.6	W74) per 100,000 age-adjusted population.
6-5	Injury by firearms	26.6	45.4		The number of deaths by firearms (ICD-10 codes W32–W34, X72–X74, X93–X95, Y22–Y24, Y 35.0) per 100,000 age-adjusted population. Causes of death attributable to firearm mortality include accidental discharge of firearms, intentional self-harm (suicide) by discharge of firearms, assault (homicide) by discharge of firearms, discharge of firearms -undetermined intent, and legal intervention involving firearm discharge. Deaths from injury by firearms exclude deaths due to explosives and other causes indirectly related to firearms.
6-6	Drug-induced deaths	72.7	79.1		The number of drug-induced deaths (ICD-10 codes F11.0–F11.5, F11.7–F11.9, F12.0–F12.5, F12.7–F12.9, F13.0–F13.5, F13.7–F13.9, F14.0–F14.5, F14.7–F14.9, F15.0–F15.5, F15.7–F15.9, F16.0–F16.5, F16.7–F16.9, F17.0, F17.3–F17.5, F17.7–F17.9, F18.0–F18.5, F18.7–F18.9, F19.0–F19.5, F19.7–F19.9, X40–X44, X60–X64, X85, Y10–Y14) per 100,000 age-adjusted population. Causes of death attributable to drug-induced mortality include selected codes from the ICD-10 title Mental and behavioral disorders due to psychoactive substance use, accidental poisoning by and exposure to drugs, medicaments and biological substances; intentional self-poisoning (suicide) by and exposure to drugs; assault (homicide) by drugs, medicaments and biological substances; and poisoning by and exposure to drugs, medicaments and biological substances, undetermined intent. Drug-induced causes exclude accidents, homicides, and other causes indirectly related to drug use. Also excluded are newborn deaths associated with mother's drug use.
6-7	Alcohol-induced deaths	181.0	644.9	24.3	The number of alcohol-induced deaths (ICD-10 codes F10, G31.2, G62.1, I42.6, K29.2, K70, R78.0, X45, X65, Y15) per 100,000 age-adjusted population. Causes of death attributable to alcohol-induced mortality include mental and behavioral disorders due to alcohol use, degeneration of nervous system due to alcohol, alcoholic polyneuropathy, alcoholic cardiomyopathy, alcoholic gastritis, alcoholic liver disease, finding of alcohol in blood, accidental poisoning by and exposure to alcohol, intentional self-poisoning by and exposure to alcohol, undetermined intent. Alcohol-induced causes exclude accidents, homicides, and other causes indirectly related to alcohol use.
6-8	Intentional self-harm (suicide)	39.7	104.6	19.4	100,000 age-adjusted population.
6-9	Assault (homicide)	23.7	192.6	8.1	The number of deaths due to assault (homicide; ICD-10 codes X85-X09, Y87.1) per 100,000 age-adjusted population.

7. PREMATURE MORTALITY

(Number of deaths per 100,000 persons in specific group)

Figure Number	Indicator	2021 rate for American Indians	Percent above or below the average rate for all groups	Average rate for all groups in 2021	Definitions
7-1	Mortality of children 1-14 years old	50.2	153.5	19.8	The number of deaths per 100,000 children 1-14 years old.
7-2	Mortality of adolescents 15-19 years old	181.1	116.4	83.7	The number of deaths per 100,000 adolescents 15-19 years old.
7-3	Mortality of young adults 20-44 years old	1012.5	285.6	262.6	The number of deaths per 100,000 young adults 20-44 years old.
7-4	Mortality of middle-aged adults 45-64 years old	2322.8	149.2	932.1	The number of deaths per 100,000 middle-aged adults 45-64 years old.

8. AGE-SPECIFIC MORTALITY

Figure Number	Indicator	2021 rate for American Indians	Percent above or below the average for all groups*	Average rate for all groups in 2021	Definitions
8-1	Median age at death from all causes	59.0	20.3	74.0	The median age (the age in the middle) of Arizonans who died in 2021.
8-2	Median age at death from all causes among males	56.0	22.2	72.0	The median age (the age in the middle) of Arizona males who died in 2021.
8-3	Median age at death from all causes among females	63.0	18.2	77.0	The median age (the age in the middle) of Arizona females who died in 2021.
8-4	Percent of total deaths before expected years of life reached	79.1	50.1	52.7	The number of deaths before reaching 78.8 years (the life expectancy of all U.S. residents in 2021) per 100 deaths.

Note: For consistency, the signs + and - for scores indicating the median age at death as higher or lower than the statewide average were reversed.

9. REPORTABLE DISEASES

(Number of cases per 100,000 persons in specific group)

Figure Number	Indicator	2021 rate for American Indians	Percent above or below the average rate for all groups	Average rate for all groups in 2021	Definitions
9-1	Incidence of early syphilis	171.6	262.0	47.4	Number of reported cases of early syphilis per 100,000 population.
9-2	Incidence of gonorrhea	412.4	62.9	253.2	Number of reported cases of gonorrhea per 100,000 population.
9-3	Incidence of chlamydia	791.8	38.9	569.9	Number of reported cases of chlamydia per 100,000 population.
9-4	Incidence of HIV/AIDS	14.5	30.6	11.1	Number of reported cases of HIV/AIDS per 100,000 population.