

**TABLE 6-6
SELECTED SUBSTANCES INVOLVED IN UNDETERMINED DRUG POISONING DEATHS, ARIZONA, 2005**

| | Total | Gender | | Race/ethnicity | | | | Age group | | | | Marital status | | | | |
|---|-------|--------|----|--------------------|--------------------|---------------------------|----------------------------------|-----------|-------|-------|-----|----------------|---------|---------|----------|------------|
| | | M | F | White non-Hispanic | Hispanic or Latino | Black or African American | American Indian or Alaska Native | 15-19 | 20-44 | 45-64 | 65+ | Never married | Married | Widowed | Divorced | Not stated |
| Heroin | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other opioids | 15 | 5 | 10 | 14 | 0 | 1 | 0 | 1 | 8 | 6 | 0 | 5 | 5 | 0 | 4 | 1 |
| Methadone | 5 | 1 | 4 | 5 | 0 | 0 | 0 | 0 | 2 | 3 | 0 | 2 | 1 | 1 | 1 | 0 |
| Other synthetic narcotics | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 0 |
| Cocaine | 4 | 4 | 0 | 2 | 0 | 2 | 0 | 0 | 3 | 1 | 0 | 4 | 0 | 0 | 0 | 0 |
| Other and unspecified narcotics | 8 | 2 | 6 | 7 | 0 | 0 | 1 | 0 | 4 | 4 | 0 | 2 | 0 | 2 | 2 | 2 |
| Tricyclic and tetracyclic antidepressants | 4 | 0 | 4 | 4 | 0 | 0 | 0 | 0 | 1 | 3 | 0 | 0 | 1 | 1 | 2 | 0 |
| Other/unspecified antidepressants | 3 | 2 | 1 | 2 | 1 | 0 | 0 | 0 | 2 | 1 | 0 | 2 | 0 | 1 | 0 | 0 |
| Psychostimulants (e.g., methamphetamine) | 7 | 3 | 4 | 5 | 1 | 0 | 1 | 1 | 3 | 3 | 0 | 2 | 2 | 0 | 2 | 1 |
| Other/unspecified psychotropic drugs | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Barbiturates | 3 | 1 | 2 | 2 | 1 | 0 | 0 | 0 | 1 | 1 | 1 | 1 | 1 | 0 | 1 | 0 |
| Benzodiazepines | 6 | 2 | 4 | 5 | 1 | 0 | 0 | 0 | 3 | 2 | 1 | 2 | 2 | 1 | 1 | 0 |
| Other antiepileptic and sedative-hypnotic drugs | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Nonopioid analgesics | 7 | 2 | 5 | 5 | 1 | 1 | 0 | 0 | 4 | 1 | 2 | 1 | 2 | 2 | 2 | 0 |
| Other and unspecified drugs | 67 | 25 | 42 | 58 | 4 | 1 | 4 | 3 | 28 | 33 | 3 | 22 | 19 | 7 | 18 | 1 |

Note: The specific substances in the above table are identified using ICD-T-codes. The sum of all identified T-codes for substances, or a specific class of substances is greater than the number of drug poisonings of undetermined intent (86).