TABLE 3-13
SUICIDE DEATHS BY GENDER, MEANS OF INJURY AND YEAR, ARIZONA RESIDENTS, 1999-2009

| | | | 1999 | 2000 | *2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|--------|---|---------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Male | Total | Count | 613 | 604 | 485 | 692 | 647 | 674 | 723 | 743 | 773 | 737 | 792 |
| | | Percent | 100% | 100% | 100% | 100% | 100% | 100% | 100% | 100% | 100% | 100% | 100% |
| | Firearm discharge or explosive material | Count | 430 | 432 | 311 | 466 | 414 | 437 | 445 | 470 | 481 | 441 | 492 |
| | | Percent | 70.1% | 71.5% | 64.1% | 67.3% | 64.0% | 64.8% | 61.5% | 63.3% | 62.2% | 59.8% | 62.1% |
| | Hanging, strangulation or suffocation | Count | 84 | 92 | 95 | 122 | 129 | 127 | 169 | 153 | 144 | 158 | 160 |
| | | Percent | 13.7% | 15.2% | 19.6% | 17.6% | 19.9% | 18.8% | 23.4% | 20.6% | 18.6% | 21.4% | 20.2% |
| | Poisoning by drugs, medicaments and biological substances | Count | 60 | 21 | 35 | 47 | 40 | 51 | 49 | 52 | 78 | 72 | 89 |
| | | Percent | 9.8% | 3.5% | 7.2% | 6.8% | 6.2% | 7.6% | 6.8% | 7.0% | 10.1% | 9.8% | 11.2% |
| | Poisoning by gases or vapors | Count | 21 | 31 | 19 | 29 | 25 | 24 | 23 | 23 | 24 | 26 | 22 |
| | | Percent | 3.4% | 5.1% | 3.9% | 4.2% | 3.9% | 3.6% | 3.2% | 3.1% | 3.1% | 3.5% | 2.8% |
| | Sharp or blunt object | Count | 9 | 9 | 10 | 8 | 8 | 15 | 16 | 24 | 16 | 13 | 17 |
| | | Percent | 1.5% | 1.5% | 2.1% | 1.2% | 1.2% | 2.2% | 2.2% | 3.2% | 2.1% | 1.8% | 2.1% |
| | Drowning and submersion | Count | 0 | 2 | 1 | 1 | 3 | 2 | 1 | 0 | 4 | 5 | 7 |
| | | Percent | 0.0% | 0.3% | 0.2% | 0.1% | 0.5% | 0.3% | 0.1% | 0.0% | 0.5% | 0.7% | 0.9% |
| | Other specified and unspecified means | Count | 9 | 17 | 14 | 19 | 24 | 18 | 21 | 21 | 26 | 22 | 5 |
| | | Percent | 1.5% | 2.8% | 2.9% | 2.7% | 3.7% | 2.7% | 2.9% | 2.8% | 3.4% | 3.0% | 0.6% |
| Female | Total | Count | 160 | 133 | 115 | 163 | 160 | 180 | 192 | 205 | 213 | 231 | 268 |
| | | Percent | 100% | 100% | 100% | 100% | 100% | 100% | 100% | 100% | 100% | 100% | 100% |
| | Firearm discharge or explosive material | Count | 65 | 54 | 47 | 78 | 62 | 61 | 63 | 84 | 60 | 89 | 104 |
| | | Percent | 40.6% | 40.6% | 40.9% | 47.9% | 38.8% | 33.9% | 32.8% | 41.0% | 28.2% | 38.5% | 38.8% |
| | Hanging, strangulation or suffocation | Count | 30 | 20 | 13 | 17 | 30 | 30 | 44 | 32 | 30 | 30 | 39 |
| | | Percent | 18.8% | 15.0% | 11.3% | 10.4% | 18.8% | 16.7% | 22.9% | 15.6% | 14.1% | 13.0% | 14.6% |
| | Poisoning by drugs, medicaments and biological substances | Count | 45 | 38 | 41 | 53 | 53 | 64 | 59 | 70 | 95 | 91 | 95 |
| | | Percent | 28.1% | 28.6% | 35.7% | 32.5% | 33.1% | 35.6% | 30.7% | 34.1% | 44.6% | 39.4% | 35.4% |
| | Poisoning by gases or vapors | Count | 9 | 10 | 8 | 8 | 9 | 10 | 9 | 12 | 13 | 11 | 18 |
| | | Percent | 5.6% | 7.5% | 7.0% | 4.9% | 5.6% | 5.6% | 4.7% | 5.9% | 6.1% | 4.8% | 6.7% |
| | · · · | Count | 2 | 0 | 0 | 0 | 0 | 4 | 5 | 1 | 4 | 0 | 5 |
| | | Percent | 1.3% | 0.0% | 0.0% | 0.0% | 0.0% | 2.2% | 2.6% | 0.5% | 1.9% | 0.0% | 1.9% |
| | Drowning and submersion | Count | 2 | 1 | 0 | 4 | 1 | 5 | 4 | 1 | 3 | 2 | 3 |
| | | Percent | 1.3% | 0.8% | 0.0% | 2.5% | 0.6% | 2.8% | 2.1% | 0.5% | 1.4% | 0.9% | 1.1% |
| | Other specified and unspecified | Count | 7 | 10 | 6 | 3 | 5 | 6 | 8 | 5 | 8 | 8 | 4 |
| | means | Percent | 4.4% | 7.5% | 5.2% | 1.8% | 3.1% | 3.3% | 4.2% | 2.4% | 3.8% | 3.5% | 1.5% |

Note: The cause-of-death titles are according to the Tenth Revision of the International Classification of Diseases (ICD-10). The causes of death for 2000-2008 are classified by ICD-10, replacing the Ninth Revision (ICD-9) used during 1979-1999. Modified counts for 2001 are not available. See footnote to Table 3-1.