

TABLE 3B-4
RATES^a OF REPORTED CASES OF GONORRHEA, CHLAMYDIA, AND EARLY SYPHILIS BY AGE AND GENDER,
ARIZONA, 2013

Age group	GONORRHEA			CHLAMYDIA			EARLY SYPHILIS		
	Males	Females	Total	Males	Females	Total	Males	Females	Total
0-4	0.0	**	**	**	**	**	0.0	0.0	0.0
5-9	0.0	**	**	0.0	0.0	0.0	0.0	0.0	0.0
10-14	3.9	9.4	6.6	11.6	60.4	35.5	0.0	0.0	0.0
15-19	178.4	321.0	247.5	656.6	2840.6	1716.2	6.5	2.7	4.7
20-24	422.1	465.4	443.0	1302.9	3832.5	2523.4	35.5	5.6	21.0
25-29	275.3	287.0	281.0	774.2	1735.9	1236.2	34.0	10.4	22.7
30-34	177.5	161.3	169.6	439.4	800.7	615.9	31.1	3.7	17.7
35-39	127.8	90.6	109.4	214.9	368.8	291.0	14.5	**	8.3
40-44	97.6	43.0	70.5	135.3	192.7	163.8	21.7	**	11.9
45-49	74.7	28.0	51.3	95.3	93.8	94.6	23.1	**	12.3
50-54	57.8	10.5	33.8	47.4	48.1	47.8	15.2	**	7.7
55-59	26.3	7.1	16.3	23.7	19.8	21.7	9.3	**	4.9
60-64	19.4	**	10.0	12.6	6.1	9.2	4.0	0.0	1.9
65-over	3.0	**	1.6	3.6	2.4	2.9	**	0.0	**
Total	101.9	94.5	98.2	264.0	667.0	466.6	13.2	1.9	7.5

Notes:^a Number of cases per 100,000 population; table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2013; denominators for unknown or transgender category are not available; rates per 100,000 population.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.