

TABLE 3B-5
FREQUENCY OF REPORTED CASES, PERCENT DISTRIBUTION AND RATES OF EARLY AND LATE SYPHILIS,
GONORRHEA, AND CHLAMYDIA BY RACE/ETHNICITY, ARIZONA, 2013

Race/ethnicity	SYPHILIS						GONORRHEA						CHLAMYDIA		
	Early			Late			Resistant			Total					
	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate
White Non-Hispanic	202	40.6	5.3	100	22.0	2.6	0	0.0	0.0	1,430	22.0	37.4	5,703	18.4	149.3
Black or African American	59	11.8	19.9	59	13.0	19.9	0	0.0	0.0	1,040	16.0	351.0	2,515	8.1	848.9
Hispanic or Latino	186	37.3	9.3	177	38.9	8.8	0	0.0	0.0	1,859	28.6	92.7	10,386	33.6	518.0
Asian or Pacific Islander	6	1.2	2.7	12	2.6	5.3	0	0.0	0.0	52	0.8	23.2	290	0.9	129.3
American Indian or Alaska Native	27	5.4	9.6	22	4.8	7.9	0	0.0	0.0	629	9.7	224.6	3,244	10.5	1158.2
Not Specified	18	3.6	N/A	85	18.7	N/A	0	0.0	N/A	1,495	23.0	N/A	8,785	28.4	N/A
Total	498	100.0	7.5	455	100.0	6.9	0	0.0	0.0	6,505	100.0	98.2	30,923	100.0	466.6

Notes: Table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2013; rates per 100,000 population.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.