

5, 2B-6, 2B-9, 2B-9, 2B-10, 2C-8, 2C-9, 2C-11, 2C-12, 2C-13, 2C-15, 2C-16, 2C-17, 2C-19, 2C-20, 2C-21, 2C-23, 2C-27, 2C-28, 2D-3, 2D-4, 5E-11, 5E-12, 5E-13, 5E-14, 5E-25, 5E-26, 5E-27, 5E-28, 5E-29, 5E-30, 5E-31, 5E-32, 5E-33, 6B-3-18
 Overexertion 2B-9
 Poisoning by drugs 2B-9
 Poisoning by other substances, gases and vapors 2B-9
 Railway accidents 2B-9
 Spider bite 2B-9
 Storms and floods 2B-9
 Suffocation and other threats to breathing 2B-9
 Venomous snakes and lizards 2B-9
 Water transport accidents 2B-9

Injuries, unintentional (accidents),
 emergency room visits for, 4D-2, 7D-2
 Cutting or piercing instruments 4D-2, 7D-2
 Drowning/submersion, 4D-2, 7D-2
 Falls, 4D-2, 7D-2
 Fire and flames, 4D-2, 7D-2
 Firearm missile, 4D-2, 7D-2
 Foreign body entering eye or other orifice, 4D-2, 7D-2
 Machinery, 4D-2, 7D-2
 Motor vehicle accident, 4D-2, 7D-2
 Natural and environmental factors, 4D-2, 7D-2
 Overexertion, 4D-2, 7D-2
 Pedal cycle, 4D-2, 7D-2
 Poisoning by drugs, 4D-2, 7D-2
 Suffocation, 4D-2, 7D-2

Injuries, unintentional (accidents), inpatient discharges with, 4D-1, 7D-1
 Falls, 4D-1, 7D-1
 Fire and flames, 4D-1, 7D-1
 Machinery, 4D-1, 7D-1
 Motor vehicle accident, 4D-1, 7D-1
 Natural and environmental factors, 4D-1, 7D-1
 Poisoning by drugs, 4D-1, 7D-1

Injury and poisoning, emergency room visits for, 4C-1, 4D-2, 7C-1, 7D-2

Inpatient deaths 2A-4, 2B-11, 5E-8

Intentional self-harm, see **Suicide**

Intrauterine hypoxia and birth asphyxia
 2C-5, 2C-6, 2C-7, 5E-20

Invertebral disc disorders, inpatient discharges for, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

Ischemic heart disease, deaths from 2B-6, 2C-27, 5E-12

J

Japanese mothers, number of births to, 1B-22

K

Knee replacement, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

L

Leading causes of death, (see also, Deaths),
 2B-1, 2B-2, 2B-3, 2B-4, 2B-5, 2B-6, 2C-28, 5E-11, 5E-12, 5E-13, 5E-14
 Age adjusted rates 2B-2, 2B-3, 2B-4, 2B-5, 5E-11
 Average/median age at death 2D-1, 2D-3, 5E-11
 County of residence 5E-11, 5E-12, 5E-13, 5E-14
 Crude death rates 2B-6, 5E-13
 Gender differences 2B-2, 2B-3, 2B-4, 2B-5, 2B-6, 2D-3, 2D-4
 Premature mortality 2D-2, 2D-4
 Race/ethnicity 2B-3, 2B-4, 2D-3, 2D-4
 Urban/rural differences 2B-5, 2D-3, 2D-4

Leading causes of death among

infants <1 year 2C-5, 2C-6, 5E-19

children 1-14 years 2C-7, 2C-8, 2C-9, 2C-10, 5E-24, 5E-25

adolescents 15-19 years 2C-11, 2C-12, 2C-13, 2C-14, 5E-26, 5E-27

young adults 20-44 years 2C-15, 2C-16, 2C-17, 2C-18, 5E-28, 5E-29

middle-aged adults 45-64 2C-19, 2C-20, 2C-21, 2C-22, 5E-30, 5E-31

elderly 65 years and older 2C-23, 2C-24, 2C-25, 2C-26, 2C-27, 5E-32, 5E-33

Selected 100+ causes 2B-6, 2C-27, 5E-11
Legal intervention, deaths due to 2B-6, 2C-27, 5E-12

Legionellosis 3A-1, 3A-2, 5F-1, 5F-2

Length of stay (average) for inpatient discharges, 4A-5

Leukemia, deaths from 2B-6, 2C-27, 5E-12

Listeriosis 3A-1, 3A-2, 5F-1, 5F-2

Live births, (see Births)

Liver disease and cirrhosis, chronic, deaths from 2B-3, 2B-4, 2B-5, 2B-6, 2C-27, 5E-11, 5E-12, 5E-13, 5E-14

Low-birthweight (LBW) births 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15, 1B-16, 1B-25, 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-34, 5B-16, 6A-1-7, 5B-17, 5B-18, 5B-19, 5B-20, 5B-21, 5B-22, 5B-23, 5B-24, 6B-1, 9A

Lung disease, chronic or acute, (see also **Medical risk factors for pregnancy**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25

Lyme disease 3A-1, 5F-1, 5F-2

Lysis of peritoneal adhesions, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

M

Malaria, deaths from, 2B-6, 2C-27, 5E-12

Malaria, reported cases, 3A-1, 3A-2, 5F-1, 5F-2

Malformed genitalia, (see also Congenital anomalies) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 1B-35, 5B-29

Malignant neoplasm, (see Cancer)

Malnutrition, deaths from 2B-6, 2C-27, 5E-12

Manic-depressive disorders, emergency room visits for, 4C-1, 7C-1

Manic-depressive disorders, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

Marital status of mother 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15, 1B-23, 1B-25, 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-24, 5B-14, 5B-15

Marriages 5G-1, 5G-2, 5G-3

Maternal deaths 1C-1, 1C-2

Measles

Number of reported cases 3A-1, 6A-2, 5F-1, 5F-2, 6B-2

Number of deaths 2B-6, 2C-28, 3A-2, 5E-12

Meconium aspiration syndrome, (see also

Abnormal conditions of the newborn) 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-34, 5B-28

Meconium, moderate/heavy, (see also

Complications of labor/delivery) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-27

Medical risk factors for pregnancy 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15, 1B-25, 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25, 5B-30

Meningitis 2B-6, 2C-28, 3A-1, 3A-2, 5E-12, 5F-1, 5F-2
Meningococcal infection 2B-6, 2C-28, 3A-1, 3A-2, 6A-2, 5E-12, 5F-1, 5F-2, 6B-2

Mental disorders, emergency room visits for, 4C-1,