

Pertussis (whooping cough)

Reported cases, 3A-1, 5F-1, 5F-2
Deaths from 2B-6, 2C-27, 2C-28, 3A-2, 5E-12

Pharyngitis, acute, emergency room visits for, 4C-2, 7C-2

Placenta previa, (*see also* **Complications of labor/delivery**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-27

Place of birth/delivery 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15

Place of death 2A-4, 2B-10, 5E-8

Plague 3A-1, 3A-2, 5F-1, 5F-2

Plurality (*see* **Multiple births**)

Pneumoconioses and chemical effects deaths from 2B-6, 2C-28, 5E-12

Pneumonia deaths from 2B-6, 2C-27, 5E-12

Pneumonia, emergency room visits for, 4C-1, 7C-1

Pneumonia, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

Pneumonitis due to solids and liquids, deaths from 2B-6, 2C-28, 5E-12

Poisoning, accidental, death from, 2B-6, 2B-9, 6A-4, 6B-4

Poisoning, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

Poisonings, emergency room visits for, 4C-1, 7C-1

Poliomyelitis, acute, deaths from 2B-6, 2C-28, 2C-29, 5E-12

Polydactyly/Syndactyly/Adactyly, (*see also*

Congenital anomalies) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 1B-35, 5B-29

Population denominators 10A-1, 10A-2, 10B-1, 10B-2, 10C-1

Postneonatal deaths 2C-1, 2C-3, 2C-4, 5E-20, 5E-21

Precipitous labor, (*see also* **Complications of labor/delivery**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-27

Pregnancies

Among girls 14 and younger 1A-7, 5A-4

Among girls 17 and younger 1A-2, 5A-3

Among girls 15-17 1A-3, 5A-4

Among girls 15-19 1A-7, 5A-3

Among girls 18-19 1A-4, 5A-4

Among girls 19 and younger 1A-5, 1A-7, 5A-2

By county of residence 5A-1, 5A-2, 5A-3, 5A-4

By outcome 1A-1, 1A-2, 1A-3, 1A-4, 1A-5, 1A-6, 1A-7, 5A-1, 5A-4

By race/ethnicity 1A-1, 1A-2, 1A-3, 1A-4, 1A-5, 1A-6

By women's age 1A-6, 1A-7, 5A-4

Pregnancy, childbirth and the puerperium, deaths related to 1C-2, 2B-6, 2C-27, 5E-12

Pregnancy with abortive outcome, deaths from 1C-2, 2B-6, 2C-28, 5E-12

Premature mortality 2D-2, 2D-4

Prenatal care

Average number of visits 5B-22

Number of visits 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15, 1B-25, 1B-26, 1B-27, 1B-28, 1B-31, 1B-32, 1B-33, 1B-34, 5B-12, 5B-21, 9A

Trimester of entry 1B-1, 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15, 1B-25, 1B-26, 1B-27, 1B-28, 1B-30, 1B-32, 1B-33, 1B-34, 5B-11, 9A

Preterm births 1B-2, 1B-3, 1B-4, 1B-5, 1B-6, 1B-7, 1B-8, 1B-9, 1B-10, 1B-11, 1B-12, 1B-13, 1B-14, 1B-15, 1B-25, 1B-26, 1B-27, 1B-28, 1B-32, 1B-33, 1B-34, 5B-22, 6A-1, 5B-23, 5B-24, 5B-30, 6B-2-1

Previous infant 4,000+ grams (*see also*

Medical risk factors for pregnancy) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25

Previous preterm or small for gestational age (SGA) infant (*see also* **Medical risk factors for pregnancy**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25

Prolonged labor, (*see also* **Complications of labor/delivery**) 1B-26, 1B-28, B-32, 1B-33, 1B-34, 5B-27

Prostate cancer, deaths from 2B-2, 2B-3, 2B-4, 2B-5, 2B-6, 2C-27, 2C-28, 2D-3, 2D-4, 5E-11, 5E-12, 5E-13, 5E-14, 5E-32, 5E-33

Prostatectomy, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

Psychoses, inpatient discharges with 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

Puerto-Rican mothers, number of births to, 1B-22

R

Rabies 3A-1, 3A-2, 5F-1, 5F-2

Rates

Age-adjusted death rates 2B-2, 2B-3, 2B-4, 2B-5, 5E-5, 5E-11

Age-specific death rates 2C-1, 2C-2, 2C-3, 2C-6, 2C-7, 2C-8, 2C-9, 2C-11, 2C-12, 2C-13, 2C-15, 2C-16, 2C-17, 2C-19, 2C-20, 2C-21, 2C-23, 2C-24, 2C-25, 2C-28, 5E-25, 5E-27, 5E-29, 5E-31, 5E-33,

Birth rates 1A-1, 1A-2, 1A-3, 1A-4, 1A-5, 1A-6, 1B-1, 5B-2

Cause-specific death rates 2B-2, 2B-3, 2B-4, 2B-5, 2B-6, 2C-6, 2C-7, 2C-8, 2C-9, 2C-11, 2C-12, 2C-13, 2C-15, 2C-16, 2C-17, 2C-19, 2C-20, 2C-21, 2C-23, 2C-24, 2C-25, 5E-11, 5E-13, 5E-19, 5E-25, 5E-27, 5E-29, 5E-31, 5E-33

Crude death rates 2B-6, 5E-13

Fertility rates 1A-1, 1A-2, 1A-3, 1A-4, 1A-5, 1A-6, 1B-1, 5A-1

Natural increase 5B-1

Pregnancy rates 1A-1, 1A-2, 1A-3, 1A-4, 1A-5, 1A-6, 1A-7, 5A-1, 5A-2, 5A-3

Rectal atresia/stenosis, (*see also* **Complications of labor/delivery**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-27

Relapsing fever, Tick 3A-1, 5F-1, 5F-2

Renal agenesis, (*see also* **Complications of labor/delivery**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-27

Renal disease, (*see also* **Medical risk factors for pregnancy**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-25

Renal failure, deaths from 2B-6, 2C-28, 5E-12

Reportable diseases (*see* **Diseases, Reportable**)

Respiratory Distress Syndrome, deaths from, (*see also* **Certain conditions originating in the perinatal period**) 2C-5, 2C-6, , 5E-19

Respiratory infection, acute upper, emergency room visits for, 4C-2, 7C-2

Respiratory system, diseases of, emergency room visits for, 4C-1, 7C-1

Respiratory system, diseases of, inpatient discharges with, 4A-1, 4A-2, 4A-3, 4A-5, 7A-1

Respiratory therapy, (inpatient procedure), 4B-1, 4B-2, 4B-3, 4B-4, 7B-1

Reyes syndrome 3A-1, 3A-2, 5F-1, 5F-2

Rheumatism, excluding back, emergency room visits for, 4C-2, 7C-2

Rh sensitization, (*see also* **Medical risk factors for pregnancy**) 1B-26, 1B-28, 1B-32, 1B-34, 5B-25

Rocky Mountain spotted fever 3A-1, 3A-2, 5F-1, 5F-2

Rubella 3A-1, 3A-2, 6A-1, 5F-1, 5F-2

Rubella syndrome, congenital 3A-1, 5F-1, 5F-2

Rupture of membranes, (*see also* **Complications of labor/delivery**) 1B-26, 1B-28, 1B-32, 1B-33, 1B-34, 5B-27

Rural areas 1B-5, 1B-7, 1B-25, 1B-26, 2B-5, 2C-1, 2C-8, 2C-12, 2C-16, 2C-20, 2C-24, 2D-3, 2D-4,