

2012 RISK PROFILE OF AMERICAN INDIANS

(Percent above or below the average
rate or ratio of all Arizona residents in
2012)

4-8.	Sudden Infant Death Syndrome (SIDS)
4-9.	Mortality of infants from low birthweight
5-7.	Lung cancer
5-17.	Alzheimer's disease
5-5.	Chronic lower respiratory diseases
5-8.	Breast cancer
4-1.	Incidence of hyaline membrane disease
5-10.	Colorectal cancer
2-2.	Tobacco use during pregnancy
5-11.	Prostate cancer
5-6.	Malignant neoplasm (cancer)
2-9.	Genital herpes during pregnancy
5-3.	Coronary heart disease
6-6.	Drug-induced deaths
5-2.	Diseases of the heart
4-2.	Incidence of newborn intensive care
5-1.	Cardiovascular disease
5-4.	Cerebrovascular disease
6-5.	Injury by firearms
4-5.	Born too small: <1,500 grams
4-4.	Born too small: <2,500 grams
4-10.	Neonatal mortality
1-2.	Pregnancies among females 14 and younger
4-3.	Born too early (preterm or < 37 weeks)
6-8.	Intentional self-harm (suicide)
4-6.	Incidence of congenital anomalies
6-3.	Fall-related injuries
5-20.	Total mortality from all causes
8-3.	Median age at death among females
4-7.	Infant mortality
8-1.	Median age at death from all causes
5-15.	HIV disease
8-2.	Median age at death among males
2-10.	Weight gain of less than 21 pounds
5-9.	Cervical cancer
5-19.	Essential (primary) hypertension
8-4.	Deaths before expected years of life reached
2-8.	Pregnancy-associated hypertension
7-4.	Mortality of middle-aged adults 45-64 years
1-3.	Pregnancies among females 15-17 years
9-5.	Incidence of HIV/AIDS
1-1.	Pregnancies among females 19 and younger
2-4.	Chronic hypertension
1-4.	Pregnancies among females 18-19 years
2-5.	Dysfunctional labor
3-3.	No prenatal care
9-1.	Incidence of early syphilis
2-1.	Unmarried mothers
9-4.	Incidence of genital herpes
2-6.	Anemia during pregnancy
3-1.	No care in the 1st trimester
4-11.	Postneonatal mortality
2-7.	Diabetes during pregnancy
9-2.	Incidence of gonorrhea
6-1.	Unintentional injuries
6-4.	Drowning mortality
7-2.	Mortality of adolescents 15-19 years
3-2.	Fewer than five prenatal visits
9-3.	Incidence of chlamydia
5-18.	Nephritis, nephrotic syndrome and nephrosis
7-3.	Mortality of young adults 20-44 years
6-9.	Assault (homicide)
7-1.	Mortality of children 1-14 years
2-3.	Alcohol use during pregnancy
6-2.	Motor vehicle-related injuries
5-12.	Influenza and pneumonia
5-16.	Septicemia
5-13.	Diabetes
5-14.	Chronic liver disease and cirrhosis
6-7.	Alcohol-induced deaths

