

## 2013 RISK PROFILE OF AMERICAN INDIANS

(Percent above or below the average rate or ratio of all Arizona residents in 2013)

5-7	Lung cancer	-72.5%
5-5	Chronic lower respiratory diseases	-65.2%
5-17	Alzheimer's disease	-51.8%
4-1	Hyaline membrane disease	-50.0%
8-4	Percent of total deaths before expected years of life reached	-40.5%
5-8	Breast cancer	-36.7%
2-9	Genital herpes during pregnancy	-34.1%
2-2	Tobacco use	-28.6%
4-9	Mortality of infants from low birthweight/short gestation	-22.6%
4-2	Newborn intensive care	-21.0%
5-6	Malignant neoplasms (cancer)	-17.0%
5-4	Cerebrovascular disease	-16.9%
5-10	Colorectal cancer	-16.7%
5-9	Cervical cancer	-15.7%
5-3	Coronary heart disease	-14.9%
6-3	Fall-related injuries	-14.1%
5-2	Diseases of the heart	-12.5%
5-1	Cardiovascular diseases	-9.1%
4-5	Born too small: very low-birthweight (<1,500 grams)	-7.8%
6-5	Injury by firearms	0.0%
4-8	Sudden Infant Death Syndrome (SIDS)	4.3%
4-4	Born too small: low-birthweight (<2,500 grams)	7.8%
4-3	Born too early (preterm or < 37 weeks)	8.8%
4-10	Neonatal mortality	9.3%
5-11	Prostate cancer	10.7%
6-6	Drug-induced deaths	14.3%
4-6	Incidence of congenital anomalies (birth defects)	17.3%
8-3	Median age at death from all causes among females	18.2%
6-4	Drowning mortality	22.1%
8-1	Median age at death from all causes	24.5%
4-7	Infant mortality	25.7%
8-2	Median age at death from all causes among males	27.3%
2-4	Chronic hypertension	28.0%
9-1	Incidence of early syphilis	28.8%
6-8	Intentional self-harm (suicide)	29.0%
5-20	Total mortality from all causes	30.9%
5-19	Essential (primary) hypertension	31.5%
2-10	Weight gain of less than 21 pounds	32.1%
2-8	Pregnancy associated hypertension	35.7%
5-15	HIV disease	45.9%
9-4	Incidence of HIV/AIDS	48.9%
2-6	Anemia during pregnancy	50.0%
1-2	Pregnancies among females 14 and younger	51.2%
1-4	Pregnancies among females 18-19 years old	52.0%
7-4	Mortality among middle-age adults 45-64 years old	52.6%
4-11	Postneonatal mortality	52.9%
2-5	Dysfunctional labor	54.7%
1-1	Pregnancies among females 19 and younger	67.4%
3-1	No care in the 1st trimester	71.4%
3-3	No prenatal care	75.0%
1-3	Pregnancies among females 15-17 years old	75.6%
2-1	Unmarried mothers	94.7%
2-7	Diabetes during pregnancy	125.7%
6-1	Unintentional injuries	126.6%
7-2	Mortality among adolescents 15-19 years old	128.7%
9-2	Incidence of gonorrhea	133.3%
3-2	Fewer than five prenatal visits	137.5%
5-16	Septicemia	147.0%
5-12	Influenza and pneumonia	148.2%
9-3	Incidence of chlamydia	148.5%
7-1	Mortality among children 1-14 years old	172.3%
7-3	Mortality among young adults 20-44 years old	175.0%
2-3	Alcohol use during pregnancy	178.1%
6-2	Motor vehicle-related injuries	178.4%
5-13	Diabetes	198.1%
5-18	Nephritis, nephrotic syndrome and nephrosis (kidney disease)	200.0%
6-9	Assault (homicide)	327.6%
5-14	Chronic liver disease and cirrhosis	439.5%
6-7	Alcohol-induced deaths	

