

**AMERICAN INDIANS' STANDING IN ARIZONA IN 2013
ON 69 INDICATORS OF HEALTH STATUS (continued)**

7. PREMATURE MORTALITY

(Number of deaths per 100,000 persons in specific group)

| Figure Number | Indicator | 2013 rate for American Indians | Percent above or below the average rate for all groups | Average rate for all groups in 2013 | Definitions |
|---------------|---|--------------------------------|--|-------------------------------------|--|
| 7-1 | Mortality of children 1-14 years old | 49.2 | 148.5 | 19.8 | The number of deaths per 100,000 children 1-14 years old. |
| 7-2 | Mortality of adolescents 15-19 years old | 103.1 | 126.6 | 45.5 | The number of deaths per 100,000 adolescents 15-19 years old. |
| 7-3 | Mortality of young adults 20-44 years old | 390.8 | 172.3 | 143.5 | The number of deaths per 100,000 young adults 20-44 years old. |
| 7-4 | Mortality of middle-aged adults 45-64 years old | 943.5 | 52.0 | 620.7 | The number of deaths per 100,000 middle-aged adults 45-64 years old. |

8. AGE-SPECIFIC MORTALITY

| Figure Number | Indicator | 2013 rate for American Indians | Percent above or below the average for all groups* | Average rate for all groups in 2013 | Definitions |
|---------------|---|--------------------------------|--|-------------------------------------|---|
| 8-1 | Median age at death from all causes | 60.0 | 22.1 | 77.0 | The median age (the age in the middle) of Arizonans who died in 2013. |
| 8-2 | Median age at death from all causes among males | 55.0 | 25.7 | 74.0 | The median age (the age in the middle) of Arizona males who died in 2013. |
| 8-3 | Median age at death from all causes among females | 67.0 | 17.3 | 81.0 | The median age (the age in the middle) of Arizona females who died in 2013. |
| 8-4 | Percent of total deaths before expected years of life reached | 76.7 | -40.5 | 54.6 | The number of deaths before reaching 78.7 years (the life expectancy of all U.S. residents in 2011) per 100 deaths. |

Note: For consistency, the signs + and - for scores indicating the median age at death as higher or lower than the statewide average were reversed.

9. REPORTABLE DISEASES

(Number of cases per 100,000 persons in specific group)

| Figure Number | Indicator | 2013 rate for American Indians | Percent above or below the average rate for all groups | Average rate for all groups in 2013 | Definitions |
|---------------|-----------------------------|--------------------------------|--|-------------------------------------|--|
| 9-1 | Incidence of early syphilis | 9.6 | 28.0 | 7.5 | Number of reported cases of early syphilis per 100,000 population. |
| 9-2 | Incidence of gonorrhea | 224.6 | 128.7 | 98.2 | Number of reported cases of gonorrhea per 100,000 population. |
| 9-3 | Incidence of chlamydia | 1158.2 | 148.2 | 466.6 | Number of reported cases of chlamydia per 100,000 population. |
| 9-4 | Incidence of HIV/AIDS | 16.2 | 45.9 | 11.1 | Number of reported cases of HIV/AIDS per 100,000 population. |