

TABLE 2D-4
PERCENT DEATHS BEFORE EXPECTED^a YEARS OF LIFE REACHED FOR THE LEADING CAUSES OF MORTALITY BY
GENDER, RACE/ETHNICITY, AND URBAN/RURAL AREAS^b, ARIZONA, 2014

	ARIZONA			Area/gender						Race/ethnicity				
	Total	Male	Female	Total Urban	Urban Male	Urban Female	Total Rural	Rural Male	Rural Female	White Non-Hispanic	Hispanic or Latino	Black or African American	American Indian or Alaska Native	Asian or Pacific Islander
Total, all causes	54.8%	62.0%	46.7%	53.8%	61.2%	45.8%	58.2%	64.4%	50.6%	50.9%	66.7%	75.2%	77.8%	58.9%
Major cardiovascular diseases	42.8%	52.8%	31.3%	41.6%	51.6%	30.4%	47.0%	56.6%	34.6%	39.6%	53.4%	69.3%	66.3%	51.7%
Diseases of heart	43.4%	53.4%	30.7%	42.1%	52.1%	29.8%	47.9%	57.2%	34.3%	40.5%	52.9%	70.8%	68.2%	53.8%
Coronary heart disease	44.8%	54.6%	31.2%	43.0%	52.8%	29.9%	51.5%	60.3%	36.6%	42.1%	53.3%	69.9%	70.5%	54.5%
Malignant neoplasms	66.9%	67.2%	66.6%	66.4%	67.1%	65.7%	68.7%	67.4%	70.3%	64.8%	75.1%	81.9%	75.0%	76.4%
Lung cancer	69.8%	71.3%	68.0%	68.5%	70.0%	66.6%	74.8%	76.0%	73.4%	69.2%	67.5%	92.4%	96.0%	58.5%
Prostate cancer	44.2%	44.2%	0.0%	44.3%	44.3%	0.0%	43.5%	43.5%	0.0%	39.7%	69.0%	56.3%	61.1%	83.3%
Breast cancer	74.6%	100.0%	74.4%	73.5%	100.0%	73.4%	79.1%	100.0%	78.6%	72.3%	81.2%	89.2%	66.7%	100.0%
Colorectal cancer	63.0%	67.3%	58.0%	61.8%	65.8%	57.4%	67.3%	72.7%	60.5%	61.2%	70.4%	75.0%	76.5%	58.3%
Malignant melanoma of skin	73.5%	72.8%	75.0%	73.1%	71.9%	76.0%	75.0%	76.9%	72.2%	73.9%	78.6%	0.0%	0.0%	0.0%
Cerebrovascular diseases	37.1%	47.1%	29.9%	35.9%	46.0%	28.6%	41.6%	51.0%	35.5%	32.7%	52.1%	60.5%	56.9%	52.4%
Chronic lower respiratory diseases	51.3%	53.3%	49.4%	50.0%	52.1%	47.9%	56.2%	56.9%	55.5%	51.0%	48.9%	67.7%	75.0%	37.5%
Accidents (unintentional injuries)	75.2%	82.2%	63.7%	73.8%	81.2%	62.0%	79.2%	84.4%	70.1%	68.2%	89.1%	90.5%	96.5%	62.5%
Motor vehicle accidents	92.6%	95.3%	86.2%	91.9%	95.6%	83.2%	94.7%	94.9%	94.4%	87.2%	98.9%	100.0%	96.9%	100.0%
Falls	32.7%	40.7%	24.7%	31.5%	39.1%	24.1%	36.1%	43.4%	28.2%	28.3%	58.3%	25.0%	85.0%	20.0%
Accidental poisoning	99.3%	99.7%	98.6%	99.1%	99.6%	98.2%	100.0%	100.0%	100.0%	99.1%	100.0%	97.8%	100.0%	100.0%
Accidental drowning	93.6%	93.5%	93.8%	92.1%	92.0%	92.3%	100.0%	100.0%	100.0%	89.4%	100.0%	100.0%	100.0%	100.0%
Influenza and pneumonia	48.6%	53.2%	43.7%	47.9%	52.0%	43.5%	49.4%	55.1%	43.4%	44.9%	64.9%	62.5%	56.3%	33.3%
Alzheimer's disease	15.6%	20.8%	12.8%	15.7%	20.2%	13.5%	14.1%	25.5%	7.0%	14.7%	20.6%	18.4%	24.3%	15.4%
Diabetes	69.1%	74.0%	62.4%	68.6%	74.5%	60.6%	71.5%	70.9%	72.3%	65.0%	74.4%	78.0%	82.4%	59.3%
Intentional self-harm (suicide)	93.1%	92.5%	95.1%	93.7%	93.1%	95.6%	91.4%	90.7%	93.5%	91.6%	97.8%	100.0%	100.0%	100.0%
Chronic liver disease and cirrhosis	92.0%	94.1%	88.8%	91.6%	93.8%	88.2%	93.4%	94.6%	91.7%	91.3%	90.3%	90.9%	96.9%	87.5%
Nephritis, nephrotic syndrome and nephrosis	42.2%	44.8%	39.7%	40.4%	42.9%	37.9%	45.6%	48.4%	42.9%	37.7%	47.5%	55.6%	80.0%	40.0%
Septicemia	58.0%	54.4%	61.7%	56.9%	52.5%	61.3%	61.1%	59.6%	62.8%	49.6%	77.6%	88.9%	86.4%	60.0%
Assault (homicide)	99.0%	99.1%	98.5%	99.2%	99.5%	98.3%	97.1%	96.2%	100.0%	98.2%	99.1%	100.0%	100.0%	100.0%
Parkinson's disease	28.1%	32.6%	21.3%	27.2%	30.8%	21.9%	32.7%	40.0%	18.2%	28.7%	20.0%	50.0%	27.8%	40.0%
Essential (primary) hypertension and hypertensive renal disease	46.2%	54.0%	39.0%	45.7%	53.2%	38.9%	48.5%	58.7%	39.6%	41.0%	58.1%	73.9%	71.0%	33.3%
Human immunodeficiency virus (HIV) disease	98.1%	100.0%	87.5%	98.9%	100.0%	92.9%	88.9%	100.0%	50.0%	100.0%	95.8%	100.0%	80.0%	100.0%
INJURY BY FIREARMS	93.0%	92.5%	95.2%	94.2%	93.8%	95.8%	88.4%	87.7%	92.3%	90.5%	99.3%	100.0%	100.0%	100.0%
DRUG-INDUCED DEATHS	98.4%	98.6%	98.0%	98.2%	98.6%	97.4%	99.1%	98.3%	100.0%	98.1%	98.9%	100.0%	100.0%	100.0%
ALCOHOL-INDUCED DEATHS	97.1%	97.0%	97.3%	96.3%	96.3%	96.4%	99.6%	99.4%	100.0%	96.4%	96.7%	91.7%	100.0%	100.0%

Notes: ^a Expected years of life at birth for all U.S. residents (78.8 years in 2013); ^b Urban = Maricopa, Pima, Pinal, and Yuma counties.