

**TABLE 3B-3
FREQUENCY OF REPORTED CASES OF GONORRHEA, CHLAMYDIA, AND EARLY SYPHILIS
BY AGE AND GENDER, ARIZONA, 2014**

Age group	GONORRHEA				CHLAMYDIA				EARLY SYPHILIS			
	Males	Females	Unknown or Transgender	Total	Males	Females	Unknown or Transgender	Total	Males	Females	Unknown or Transgender	Total
0-4	0	*	0	0†	0	0	0	0	0	0	0	0
5-9	0	0	0	0	0	*	0	0†	0	0	0	0
10-14	*	16	0	20†	24	143	0	167	0	0	0	0
15-19	503	679	0	1,182	1,606	5,876	0	7,482	36	9	0	45
20-24	1,163	1,198	0	2,361	3,454	8,878	0	12,332	161	20	0	181
25-29	894	695	0	1,589	2,047	3,903	0	5,950	131	25	0	156
30-34	549	431	0	980	1,048	1,832	0	2,880	113	16	0	129
35-39	334	218	0	552	529	859	0	1,388	74	10	0	84
40-44	225	127	0	352	338	402	0	740	75	9	0	84
45-49	196	64	0	260	193	185	0	378	88	*	0	90†
50-54	112	36	0	148	109	105	0	214	60	*	0	60†
55-59	51	17	0	68	67	61	0	128	29	*	0	30†
60-64	37	*	0	40†	30	16	0	46	14	0	0	14
65-over	29	*	0	30†	26	16	0	42	7	0	0	7
Total	4,100†	3,490†	0	7,590†	9,471	22,280†	0	31,750†	788	90†	0	880

Notes: * Cell suppressed due to non-zero count less than 6; † Sum rounded to nearest tens unit due to non-zero addend less than 6; since 2005, the table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.