

**TABLE 2B-5
AGE-ADJUSTED* MORTALITY RATES FOR SELECTED CAUSES OF DEATH BY URBAN/RURAL AREA AND GENDER,
ARIZONA, 2000**

| | Arizona | | | Urban | | | Rural | | |
|--|---------|-------|--------|-------|-------|--------|-------|-------|--------|
| | Total | Male | Female | Total | Male | Female | Total | Male | Female |
| Total, all causes | 785.6 | 941.1 | 654.6 | 778.3 | 933.0 | 649.9 | 818.2 | 978.1 | 675.6 |
| Cardiovascular disease | 276.1 | 333.1 | 230.0 | 275.5 | 332.2 | 230.5 | 276.4 | 334.7 | 226.4 |
| Diseases of heart | 206.1 | 259.3 | 163.6 | 205.8 | 259.7 | 163.5 | 205.7 | 255.5 | 162.6 |
| Malignant neoplasm | 170.4 | 204.0 | 145.8 | 170.5 | 204.6 | 145.8 | 170.3 | 201.8 | 145.6 |
| Colorectal cancer | 17.0 | 20.5 | 14.1 | 17.1 | 20.2 | 14.6 | 16.4 | 21.7 | 11.8 |
| Lung cancer | 46.1 | 59.2 | 35.8 | 46.3 | 59.8 | 35.9 | 45.4 | 56.4 | 35.7 |
| Prostate cancer | NA | 28.4 | NA | NA | 28.5 | NA | NA | 27.9 | NA |
| Female breast cancer | NA | NA | 25.4 | NA | NA | 25.7 | NA | NA | 24.2 |
| Cerebrovascular disease | 51.7 | 50.8 | 51.8 | 51.5 | 50.3 | 51.7 | 52.1 | 52.8 | 51.7 |
| Chronic lower respiratory diseases | 47.4 | 56.2 | 41.3 | 48.2 | 56.6 | 42.6 | 43.8 | 54.6 | 35.5 |
| Accidents (unintentional injuries) | 41.1 | 58.0 | 24.9 | 38.3 | 54.9 | 22.6 | 55.2 | 74.6 | 36.2 |
| Motor vehicle accidents | 17.5 | 24.4 | 10.6 | 15.3 | 21.7 | 8.8 | 28.5 | 37.6 | 19.5 |
| Falls | 7.6 | 9.8 | 5.9 | 7.5 | 10.1 | 5.6 | 7.8 | 8.9 | 6.7 |
| Accidental drowning | 1.7 | 2.5 | .9 | 1.7 | 2.5 | 1.0 | 1.8 | 2.9 | .6 |
| Accidental poisoning | 5.0 | 8.0 | 2.1 | 5.2 | 8.1 | 2.3 | 4.1 | 7.4 | .8 |
| Influenza and pneumonia | 24.4 | 29.2 | 21.1 | 23.9 | 29.2 | 20.4 | 27.0 | 29.1 | 25.0 |
| Alzheimer's disease | 21.8 | 17.5 | 24.2 | 22.6 | 17.4 | 25.4 | 18.0 | 17.7 | 18.3 |
| Diabetes | 19.0 | 21.0 | 17.2 | 18.5 | 21.2 | 16.3 | 21.0 | 20.3 | 21.1 |
| Intentional self-harm (suicide) | 14.6 | 24.7 | 5.2 | 13.1 | 22.5 | 4.4 | 21.2 | 34.1 | 9.0 |
| Chronic liver disease and cirrhosis | 12.8 | 17.7 | 8.3 | 12.1 | 16.6 | 8.0 | 16.3 | 23.4 | 9.6 |
| Nephritis | 11.7 | 13.9 | 10.4 | 11.1 | 13.2 | 9.9 | 14.4 | 16.9 | 12.4 |
| Septicemia | 6.7 | 7.3 | 6.1 | 6.2 | 6.9 | 5.6 | 8.7 | 9.0 | 8.2 |
| Assault (homicide) | 7.6 | 11.1 | 3.8 | 7.9 | 11.7 | 3.8 | 5.9 | 8.0 | 3.7 |
| Parkinson's disease | 6.8 | 10.1 | 4.7 | 7.1 | 10.8 | 4.9 | 5.2 | 7.3 | 3.7 |
| Essential (primary) hypertension | 5.6 | 6.2 | 5.1 | 5.8 | 6.2 | 5.5 | 4.6 | 6.3 | 3.3 |
| Aortic aneurysm | 5.0 | 8.0 | 2.7 | 4.7 | 7.4 | 2.7 | 6.4 | 10.6 | 2.8 |
| Congenital malformations | 3.6 | 3.6 | 3.6 | 3.5 | 3.5 | 3.6 | 3.9 | 4.3 | 3.6 |
| HIV disease | 3.2 | 5.5 | 1.0 | 3.3 | 5.7 | .9 | 2.9 | 4.4 | 1.5 |
| INJURY BY FIREARMS | 15.3 | 26.9 | 4.3 | 14.4 | 25.8 | 3.5 | 19.1 | 30.8 | 8.1 |
| DRUG-INDUCED DEATHS | 6.8 | 9.4 | 4.1 | 7.2 | 9.9 | 4.4 | 4.8 | 7.2 | 2.3 |
| ALCOHOL-INDUCED DEATHS | 8.7 | 13.5 | 4.3 | 7.4 | 11.2 | 3.8 | 15.2 | 24.4 | 6.5 |

*Adjusted to the 2000 standard U.S. population. The causes of death for 2000 are classified by the Tenth Revision of the International Classification of Diseases (ICD-10). The rate for breast cancer is per 100,000 females. The rate for prostate cancer is per 100,000 males.