

**TABLE 6
MONITORING PROGRESS TOWARD ARIZONA AND SELECTED NATIONAL YEAR 2010 OBJECTIVES (2002 STATUS)**

Focus areas and selected objectives: (In parentheses are <i>Healthy People 2010</i> objective numbers)	2010 TARGET	RATES, RATIOS OR CASES IN 2002:															
		ARIZONA	Apache	Cochise	Coconino	Gila	Graham	Greenlee	Maricopa	Mohave	Navajo	Pima	Pinal	Santa Cruz	Yavapai	Yuma	La Paz
6A. MATERNAL, INFANT AND CHILD HEALTH																	
Reduce fetal deaths at 20 or more weeks of gestation (HP16-1a)	4.1	7.1	4.5	4.7	8.5	7.1	6.1	20.2	7.4	6.0	4.9	7.0	7.7	2.6	5.7	8.1	4.3
Reduce fetal and infant deaths during perinatal period (HP16-1b)	4.5	6.4	4.5	6.4	7.4	11.4	12.2	6.7	4.0	3.7	6.0	7.3	1.3	3.1	4.8	8.6	4.5
♦Reduce infant deaths (HP16-1c)	♦6.0	6.3	5.4	7.6	7.0	11.5	12.3	10.3	6.4	4.5	4.3	7.2	5.9	5.2	4.7	2.6	4.3
Reduce neonatal deaths (HP16-1d)	2.9	4.1	3.6	2.9	5.4	8.6	10.3	10.3	4.3	3.5	.6	4.2	4.4	2.6	2.6	2.0	4.3
Reduce postneonatal deaths (HP16-1e)	1.2	2.2	1.8	4.7	1.6	2.9	2.1	0.0	2.1	1.0	3.7	3.0	1.5	2.6	2.1	.7	0.0
Increase the proportion of pregnant women who receive prenatal care in the first trimester (HP16-6a)	90%	75.7	64.8	66.9	79.5	65.4	70.2	59.8	79.1	74.8	66.7	69.3	70.8	65.4	74.8	63.2	60.0
Reduce low birth weight (LBW) (HP16-10a)	5.0%	6.8	5.1	6.8	7.3	6.9	7.8	5.2	6.7	6.8	7.2	7.4	7.7	6.3	6.8	4.9	5.2
Reduce very low birth weight (VLBW) (HP16-10b)	0.9%	1.1	0.7	1.1	1.0	1.0	1.2	0.0	1.2	0.8	0.8	1.1	1.6	0.9	0.7	0.7	0.4
Reduce preterm births (HP16-11a)	7.6%	10.0	6.6	9.1	10.1	10.1	11.3	11.3	10.4	9.3	10.5	9.8	10.3	7.7	9.7	7.5	9.6
Increase abstinence from cigarette smoking among pregnant women (HP16-17c)	99.0%	93.7	95.5	89.1	98.4	86.2	89.1	84.5	94.2	81.3	92.8	94.3	90.2	99.6	85.8	98.9	94.3
6B. RESPONSIBLE SEXUAL BEHAVIOR																	
♦ Reduce pregnancies among adolescent females aged 15 to 17 years ^a (HP9-7)	♦ 25.0	42.2	26.2	35.8	33.5	61.0	36.0	40.0	43.4	32.4	31.9	44.7	49.9	46.2	27.4	46.5	45.5
Reduce gonorrhea infections ^b (HP25-2)	19.1	68.9	45.7	14.5	24.0	18.9	2.9	0.0	90.3	13.3	95.1	50.5	38.0	27.5	7.2	22.4	0.0
Reduce the incidence of primary and secondary syphilis ^b (HP25-3)	0.2	3.7	1.4	0.8	4.0	0.0	2.9	0.0	4.7	0.0	8.8	3.3	0.0	0.0	0.0	0.0	0.0

♦ Indicates objectives and targets identified in *Healthy Arizona 2010*.

Notes: Fetal and perinatal deaths are per 1,000 live births+fetal deaths. Perinatal period = 28 weeks of gestation to 7 days after birth. Infant, neonatal, and postneonatal deaths are per 1,000 live births. The pregnancy rates are per 1,000 females 15-17 years old. The incidence rates of gonorrhea and primary and secondary syphilis are per 100,000 population (Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV/STD/HCV Services). All other proportions and ratios are per 100 live births. Preterm births = births prior to 37 completed weeks of gestation.

The rates based on fewer than 10 cases are not statistically reliable. See Tables 5A-1, 5B-16, 5C-1, 5C-3, 5E-16, 5E-21, 5E-23, 5F-1 for the numerators.