

TABLE 6 (continued)
MONITORING PROGRESS TOWARD ARIZONA AND SELECTED NATIONAL YEAR 2010 OBJECTIVES (2002 STATUS)

Focus areas and selected objectives: (In parentheses are <i>Healthy People 2010</i> objective numbers)	2010 TARGET	RATES, RATIOS OR CASES IN 2002:															
		ARIZONA	Apache	Cochise	Coconino	Gila	Graham	Greenlee	Maricopa	Mohave	Navajo	Pima	Pinal	Santa Cruz	Yavapai	Yuma	La Paz
6C. VACCINE PREVENTABLE DISEASES																	
Reduce the rate of hepatitis A (HP14-6)	4.5	5.6	1.4	8.9	3.2	0.0	2.9	0.0	5.5	3.0	6.9	6.2	4.2	32.6	1.1	7.1	4.3
Reduce the rate of meningococcal disease (HP14-7)	1.0	0.6	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	2.0	0.3	2.1	2.5	0.0	0.6	0.0
Reduce the rate of tuberculosis (HP14-11)	1.0	4.8	14.3	1.6	7.2	0.0	2.9	0.0	5.2	2.4	8.9	2.7	7.3	5.0	1.1	8.2	0.0
6D. INJURY AND VIOLENCE																	
Reduce firearm-related deaths (HP15-3)	4.1	17.3	19.5	15.8	10.8	23.3	23.7	0.0	16.8	23.2	15.4	17.6	17.5	0.0	26.6	10.2	18.6
Reduce deaths caused by unintentional injuries (HP15-13)	17.5	45.2	112.2	50.4	57.0	62.3	38.7	61.6	40.6	44.4	104.0	46.9	52.9	32.6	50.7	39.2	104.0
♦ Reduce deaths caused by motor vehicle crashes (HP15-15)	♦ 16.0	18.5	70.4	14.9	29.1	25.2	18.6	61.6	15.5	18.6	61.0	15.9	26.8	16.2	21.8	23.4	48.9
Reduce deaths from falls (HP15-27)	3.0	8.6	12.5	10.6	10.3	2.8	3.2	0.0	9.5	4.6	12.3	7.8	5.9	2.6	11.2	3.3	12.9
♦ Reduce homicides (HP15-32)	♦ 5.0	8.6	23.8	3.4	7.3	4.7	6.9	12.6	8.6	3.3	13.2	10.4	9.1	2.7	5.4	4.0	7.7
♦ Reduce the suicide rate (HP18-1)	♦ 10.0	15.9	19.9	15.5	16.0	28.0	25.8	0.0	14.2	28.5	24.7	16.0	18.6	5.3	26.1	11.1	19.9
♦ Reduce the suicide rate among adolescents aged 15 to 19 years	♦ 10.0	9.9	0.0	0.0	0.0	28.4	30.4	10.0	30.5	42.4	10.7	0.0	0.0	0.0	0.0	0.0	9.9
♦ Reduce the suicide rate among older adults aged 65 years and older	♦ 10.0	23.2	0.0	49.2	11.4	19.1	24.7	.0	18.7	44.0	29.5	27.0	22.4	0.0	35.3	17.8	38.1
6E. CANCER																	
Reduce the overall cancer death rate (HP3-1)	159.9	168.6	118.9	176.5	149.1	206.2	160.4	137.3	168.0	207.0	136.3	173.8	175.9	160.6	169.6	136.6	110.6
Reduce the lung cancer death rate (HP3-2)	44.9	46.6	14.4	45.0	29.8	41.7	35.3	21.0	46.9	72.8	27.5	46.0	52.1	41.3	52.0	36.8	34.1
Reduce the breast cancer death rate (HP3-3)	22.3	22.5	19.9	27.1	20.7	42.0	16.5	0.0	23.9	22.6	22.5	19.4	17.9	10.1	23.7	17.6	8.5
Reduce the colorectal cancer death rate (HP3-5)	13.9	16.5	11.7	17.9	14.4	19.7	23.7	0.0	16.7	17.2	13.5	17.6	20.3	11.3	14.5	8.5	7.6
Reduce the prostate cancer death rate (HP3-7)	28.8	24.8	18.6	28.9	33.5	33.0	6.6	0.0	25.6	31.8	12.9	25.7	14.6	39.4	25.9	14.3	15.6

♦ Indicates objectives and targets identified in *Healthy Arizona 2010*.

Notes: The rates of hepatitis A, meningococcal disease, and tuberculosis are per 100,000 persons. (Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of Infectious Disease Services, Infectious Disease Epidemiology Section). Except the age-specific mortality rates for suicide, all mortality rates are adjusted to the 2000 standard population and expressed per 100,000 population. The rates based on fewer than 10 cases are not statistically reliable. See Tables 5E-12, 5E-28 and 5F-1 for the numerators.