

**TABLE 3B-5  
 FREQUENCY OF REPORTED CASES, PERCENT DISTRIBUTION AND RATES OF EARLY AND LATE SYPHILIS, GONORRHEA,  
 CHLAMYDIA AND HERPES BY ETHNICITY, ARIZONA, 2003**

Race/ Ethnicity	SYPHILIS						GONORRHEA						CHLAMYDIA			HERPES		
	Early			Late			Resistant			Total			Cases	%	Rate	Cases	%	Rate
	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate	Cases	%	Rate						
<b>White Non-Hispanic</b>	150	36	4.2	149	22	4.1	1	50	0	1,205	33	33.4	3,633	28	100.7	424	47	11.8
<b>Black or African American</b>	34	8	18.2	78	11	41.9	0	0	0	669	18	359.7	1,000	7	537.6	43	5	23.1
<b>Hispanic or Latino</b>	138	33	9.7	366	55	25.7	1	50	0.1	1,016	28	71.4	4,687	37	329.1	204	22	14.3
<b>Asian or Pacific Islander</b>	1	0	0.9	2	0	1.7	0	0	0	18	1	15.3	102	1	86.4	9	1	7.6
<b>American Indian or Alaska Native</b>	59	14	20.1	31	5	10.6	0	0	0	302	9	103.1	1,828	14	623.9	89	10	30.4
<b>Not Specified</b>	36	9	na	33	7	na	0	0	0	366	11	na	1,535	13	na	141	15	Na
<b>Total</b>	418	100	7.4	659	100	12.4	2	100	0.02	3,576	100	63.5	12,785	100	227.1	910	100	16.2

\* Quantity equal or less than 0.05

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV/STD Services