

**TABLE 2D-4
PERCENT DEATHS BEFORE EXPECTED* YEARS OF LIFE REACHED FOR THE LEADING CAUSES OF MORTALITY BY GENDER,
RACE/ETHNICITY AND URBAN/RURAL AREAS**, ARIZONA, 2004**

	TOTAL STATE			AREA/GENDER:						RACE/ETHNICITY:				
	TOTAL	MALE	FEMALE	TOTAL URBAN	URBAN MALE	URBAN FEMALE	TOTAL RURAL	RURAL MALE	RURAL FEMALE	WHITE NON-HISPANIC	HISPANIC or LATINO	BLACK or AFRICAN-AMERICAN	AMERICAN INDIAN or ALASKA NATIVE	ASIAN or PACIFIC ISLANDER
Total, all causes	53.0%	61.0%	44.1%	51.7%	59.9%	42.9%	57.9%	65.3%	49.1%	48.3%	71.5%	74.3%	80.3%	64.3%
Major cardiovascular diseases	40.6%	50.9%	30.0%	38.6%	49.0%	28.3%	48.8%	58.2%	37.8%	37.2%	57.1%	65.7%	62.3%	57.6%
Diseases of heart	42.4%	52.6%	30.9%	40.4%	50.7%	29.1%	50.6%	59.7%	38.7%	39.4%	59.6%	67.1%	58.6%	53.8%
Coronary heart disease	43.3%	53.5%	30.8%	40.9%	51.2%	28.9%	53.6%	62.6%	40.3%	40.6%	58.9%	64.8%	58.5%	53.4%
Malignant neoplasms	65.3%	66.4%	64.0%	64.8%	65.8%	63.6%	67.4%	68.6%	65.7%	63.7%	73.5%	74.4%	75.5%	88.9%
Lung cancer	68.3%	70.4%	65.6%	67.9%	69.3%	66.1%	69.9%	73.9%	64.0%	67.8%	71.1%	71.9%	76.2%	94.4%
Prostate cancer	35.7%	35.7%	0.0%	35.5%	35.5%	0.0%	36.3%	36.3%	0.0%	34.7%	29.1%	57.9%	62.5%	0.0%
Female breast cancer	73.0%	100.0%	73.0%	72.7%	100.0%	72.7%	74.4%	0.0%	74.4%	69.6%	92.5%	89.5%	94.1%	66.7%
Colorectal cancer	60.1%	65.0%	54.5%	58.2%	64.1%	52.2%	67.4%	67.8%	66.7%	58.2%	73.2%	70.4%	66.7%	85.7%
Malignant melanoma of skin	75.0%	78.8%	66.7%	72.3%	76.3%	64.1%	84.8%	87.5%	77.8%	74.3%	100.0%	0.0%	100.0%	0.0%
Cerebrovascular diseases	33.5%	44.5%	26.0%	31.9%	42.9%	24.5%	41.0%	51.9%	33.1%	29.0%	49.6%	62.1%	78.4%	72.0%
Chronic lower respiratory diseases	45.7%	48.0%	43.5%	45.9%	47.5%	44.4%	44.9%	49.8%	39.4%	45.3%	45.0%	70.8%	69.2%	44.4%
Accidents (unintentional injuries)	77.7%	84.5%	65.9%	76.2%	83.4%	63.3%	82.5%	88.6%	73.5%	71.1%	91.3%	94.4%	93.6%	76.5%
Motor vehicle accidents	92.7%	93.7%	90.7%	92.6%	93.5%	90.9%	92.7%	94.2%	90.3%	89.5%	97.2%	100.0%	96.7%	80.0%
Falls	27.8%	38.8%	18.6%	25.3%	35.0%	17.0%	40.0%	57.5%	26.0%	23.5%	47.6%	62.5%	66.7%	50.0%
Accidental poisoning	98.3%	98.9%	96.7%	98.7%	99.4%	96.7%	96.2%	95.8%	96.8%	98.2%	97.5%	100.0%	100.0%	100.0%
Accidental drowning	91.3%	93.2%	87.1%	90.7%	93.2%	85.2%	94.4%	92.9%	100.0%	87.1%	96.0%	100.0%	100.0%	100.0%
Influenza and pneumonia	35.6%	39.8%	31.2%	34.8%	39.0%	30.3%	38.7%	42.4%	34.5%	30.2%	63.3%	76.9%	63.0%	25.0%
Alzheimer's disease	10.0%	13.2%	8.5%	9.3%	12.6%	7.8%	13.7%	16.0%	12.6%	9.2%	22.2%	6.5%	25.0%	0.0%
Diabetes	61.2%	67.4%	55.0%	60.9%	67.5%	54.1%	62.1%	66.7%	57.8%	56.1%	69.0%	75.0%	81.4%	46.2%
Intentional self-harm (suicide)	92.0%	91.2%	95.0%	91.7%	90.7%	95.0%	93.2%	92.8%	95.0%	90.2%	98.3%	95.0%	100.0%	100.0%
Chronic liver disease and cirrhosis	90.4%	93.9%	83.4%	88.7%	92.6%	81.6%	95.2%	96.8%	90.5%	88.2%	90.3%	87.5%	99.0%	100.0%
Nephritis, nephrotic syndrome and nephrosis	49.8%	51.7%	47.7%	48.0%	50.6%	45.1%	55.4%	55.7%	55.1%	40.7%	66.7%	70.4%	86.7%	33.3%
Septicemia	62.9%	66.1%	59.6%	63.9%	66.9%	60.7%	60.2%	63.8%	56.7%	57.4%	74.4%	70.6%	82.4%	100.0%
Assault (homicide)	98.1%	99.0%	94.0%	98.3%	99.4%	93.0%	97.0%	96.3%	100.0%	95.6%	100.0%	97.6%	100.0%	100.0%
Parkinson's disease	25.9%	29.1%	20.5%	26.1%	29.2%	20.5%	24.6%	28.6%	20.7%	26.2%	16.7%	0.0%	60.0%	0.0%
Essential (primary) hypertension and hypertensive renal disease	36.4%	44.1%	30.4%	35.3%	41.4%	30.7%	43.4%	60.0%	28.6%	28.1%	55.6%	56.0%	77.8%	75.0%
Human immunodeficiency virus (HIV) disease	98.7%	99.3%	94.7%	98.6%	99.2%	94.1%	100.0%	100.0%	100.0%	97.8%	100.0%	100.0%	100.0%	100.0%
INJURY BY FIREARMS	92.9%	92.7%	94.4%	93.4%	93.2%	95.3%	90.6%	90.5%	90.9%	88.7%	99.2%	100.0%	100.0%	100.0%
DRUG-INDUCED DEATHS	98.0%	98.3%	97.4%	98.2%	98.5%	97.5%	96.8%	96.8%	96.8%	97.5%	99.1%	100.0%	100.0%	100.0%
ALCOHOL-INDUCED DEATHS	95.7%	96.2%	94.2%	95.5%	95.9%	94.2%	96.5%	97.2%	94.1%	93.7%	97.6%	100.0%	100.0%	100.0%

* Expected years of life at birth for all U.S. residents.

** Urban = Maricopa, Pima, Pinal and Yuma counties.

Note: The causes of death for 2004 are classified by the Tenth Revision of the International Classification of Diseases (ICD-10).