

TABLE 5E-28
RATES* FOR THE LEADING CAUSES OF DEATH AMONG ADOLESCENTS (15-19 YEARS) BY COUNTY OF RESIDENCE,
ARIZONA, 2006

| Underlying cause of death | ARIZONA | County of residence | | | | | | | | | | | | | | |
|---|---------|---------------------|---------|----------|-------|--------|----------|----------|--------|--------|------|-------|------------|---------|------|--------|
| | | Apache | Cochise | Coconino | Gila | Graham | Greenlee | Maricopa | Mohave | Navajo | Pima | Pinal | Santa Cruz | Yavapai | Yuma | La Paz |
| Accident (unintentional injury) | 43.3 | 86.8 | 9.3 | 70.5 | 25.7 | 59.5 | 0.0 | 40.4 | 25.8 | 101.4 | 39.5 | 75.7 | 24.6 | 82.9 | 0.0 | 0.0 |
| Motor vehicle accident | 34.2 | 37.2 | 9.3 | 52.9 | 0.0 | 59.5 | 0.0 | 32.5 | 17.2 | 101.4 | 28.2 | 70.3 | 24.6 | 60.3 | 0.0 | 0.0 |
| Accidental drowning and submersion | 2.0 | 12.4 | 0.0 | 8.8 | 0.0 | 0.0 | 0.0 | 2.0 | 0.0 | 0.0 | 1.4 | 0.0 | 0.0 | 7.5 | 0.0 | 0.0 |
| Accidental discharge of firearms | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Accidental poisoning | 3.6 | 12.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.9 | 0.0 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Assault (homicide) | 18.0 | 12.4 | 0.0 | 8.8 | 0.0 | 0.0 | 0.0 | 21.9 | 17.2 | 18.4 | 15.5 | 16.2 | 0.0 | 7.5 | 12.9 | 0.0 |
| By discharge of firearms | 13.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 19.2 | 0.0 | 0.0 | 14.1 | 5.4 | 0.0 | 0.0 | 6.5 | 0.0 |
| By other means | 4.1 | 12.4 | 0.0 | 8.8 | 0.0 | 0.0 | 0.0 | 2.7 | 17.2 | 18.4 | 1.4 | 10.8 | 0.0 | 7.5 | 6.5 | 0.0 |
| Intentional self-harm (suicide) | 13.0 | 24.8 | 27.8 | 17.6 | 25.7 | 29.7 | 145.3 | 11.4 | 25.8 | 46.1 | 8.5 | 10.8 | 24.6 | 0.0 | 6.5 | 0.0 |
| By discharge of firearms | 6.4 | 0.0 | 9.3 | 0.0 | 0.0 | 29.7 | 145.3 | 6.7 | 17.2 | 27.7 | 1.4 | 0.0 | 24.6 | 0.0 | 6.5 | 0.0 |
| By other means | 6.6 | 24.8 | 18.5 | 17.6 | 25.7 | 0.0 | 0.0 | 4.7 | 8.6 | 18.4 | 7.0 | 10.8 | 0.0 | 0.0 | 0.0 | 0.0 |
| Malignant neoplasms | 3.4 | 24.8 | 9.3 | 0.0 | 0.0 | 29.7 | 0.0 | 2.7 | 0.0 | 0.0 | 1.4 | 10.8 | 0.0 | 7.5 | 0.0 | 0.0 |
| Diseases of heart | 1.6 | 0.0 | 9.3 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 0.0 | 24.6 | 0.0 | 0.0 | 0.0 |
| Congenital malformations | 0.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 6.5 | 0.0 |
| Cerebrovascular diseases | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Influenza and pneumonia | 0.9 | 12.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 8.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Total, all causes | 93.8 | 198.5 | 55.5 | 123.4 | 102.9 | 118.9 | 145.3 | 91.3 | 77.4 | 184.5 | 81.8 | 124.4 | 73.9 | 113.0 | 38.8 | 0.0 |

NOTE: There were no deaths among adolescents 15-19 years in Greenlee County. The rates based on fewer than 10 deaths (see Table 5E-27) are not statistically reliable. All rates are per 100,000 adolescents 15-19 years old.