

**TABLE 3B-4
RATES¹ OF REPORTED CASES OF GONORRHEA, CHLAMYDIA, EARLY SYPHILIS AND GENITAL HERPES
BY AGE AND GENDER, ARIZONA, 2009**

Age group	GONORRHEA			CHLAMYDIA			EARLY SYPHILIS			GENITAL HERPES		
	Males	Females	Total	Males	Females	Total	Males	Females	Total	Males	Females	Total
0-4	0.8	0.4	0.6	3.1	4.9	4.0	0.0	0.0	0.0	2.4	2.9	2.6
5-9	0.0	0.0	0.0	0.4	1.7	1.1	0.0	0.0	0.0	0.4	1.3	0.9
10-14	1.7	11.4	6.4	12.5	98.4	54.4	0.0	0.0	0.0	0.4	0.4	0.4
15-19	117.6	205.4	160.1	685.3	3070.0	1838.5	7.2	2.7	5.0	6.4	27.2	16.4
20-24	227.8	240.5	233.8	1015.0	3157.8	2029.3	25.4	11.1	18.6	14.1	48.1	30.2
25-29	129.2	98.4	114.6	536.3	1223.1	863.3	21.7	8.0	15.3	15.2	29.3	21.9
30-34	92.2	49.4	71.5	278.7	558.3	413.7	19.3	1.8	10.8	13.0	31.0	21.7
35-39	55.2	32.9	44.3	143.0	271.3	205.7	16.4	4.5	10.6	11.6	20.3	15.9
40-44	38.1	12.5	25.5	86.2	116.6	101.2	23.4	4.0	13.8	12.1	18.7	15.4
45-54	32.5	5.3	18.8	36.3	41.3	38.9	12.6	1.6	7.0	9.5	15.3	12.4
55-64	7.5	1.2	4.2	10.6	10.4	10.5	5.3	0.6	2.9	8.1	8.4	8.3
65-over	1.8	0.4	1.0	3.9	3.4	3.6	0.5	0.0	0.2	3.1	3.8	3.5
Total	53.8	44.8	49.3	209.1	579.7	394.2	10.4	2.5	6.5	7.9	15.6	11.7

¹Number of cases per 100,000 population.

Note: Table includes all positive laboratory results for chlamydia and gonorrhea with or without communicable disease report in 2009.

Note: Denominators for unknown or transgender category are not available.

Rates per 100,000 population.

Source: Arizona Department of Health Services, Bureau of Epidemiology and Disease Control Services, Office of HIV / STD.